

Where the Buffalo Roam: Infinite Processes and Infinite Complexity

Stephen Semmes
Department of Mathematics
Rice University

Preface

Much attention has been paid in recent years to questions of *complexity*, how much time or memory does it take to make a certain construction, or to answer a particular question, or what is the length of the shortest path with prescribed properties, and so forth. In mathematics one allows infinite processes and infinite constructions, and the time seems to be ripe to look at measurements of their complexity.

We shall try to paint a picture through examples.

This manuscript (modulo a few modest changes) was distributed informally as IHES preprint M/96/77. The author was partially supported by the National Science Foundation.

Contents

Preface	ii
1 Introduction	1
2 Examples from the infinite world	3
2.1 Continuity	3
2.2 Uniform continuity	4
2.3 Classical Banach spaces	6
2.4 The open mapping theorem	8
2.5 Category versus measure	10
3 Symmetry and language	14
4 The existence of limits	18
4.1 The Riemann-Lebesgue lemma	18
4.2 Lebesgue's theorem	21
4.3 The ergodic theorem	24
4.4 Differentiability almost everywhere	26
5 Using infinite processes	29
5.1 Mostow rigidity	29
5.2 Groups of polynomial growth	32
5.3 A product expansion for $\sin z$	33
5.4 Transversality	34
5.5 Homeomorphisms	36
6 Finite constructions and their limits	40
6.1 The Hilbert transform	41
6.2 Topology	43

6.3	Finite sets and geometry	45
6.4	Finitely generated groups	48
6.5	Comments about symmetry and infinity	49
7	Measuring infinite complexity, I	50
7.1	Functions and polynomials	50
7.2	Polynomials, schmolynomials	55
7.3	Linear operators	60
7.4	General comments about operators	64
8	Measuring infinite complexity, II	67
8.1	Sizes of sets	67
8.2	Complexity of sets	71
8.3	Topological dimension	73
8.4	Topological dimension, size, and structure	76
8.5	Lipschitz mappings	78
8.6	Differentiability, rectifiability, etc.	80
8.7	Wild embeddings	82
8.8	The Whitehead continuum	83
8.9	Decompositions	84
8.10	Domains in Euclidean spaces	85
8.11	Spaces of bounded geometry	87
9	Some results about geometric complexity	88
9.1	Discrete parameterizations	88
9.2	A topological result about infinite complexity	91
9.3	Almost flat hypersurfaces	92
9.4	Big pieces of bilipschitz mappings	93
	Bibliography	96

Chapter 1

Introduction

In the formalization of mathematics there are obvious philosophical questions of what is real and what is mere abstraction, and how mathematics might treat the two differently. In the daily life of the working mathematician there is not much point to the broader debate; the matter is typically clear in actual practice. We know when we have given an argument which is nonconstructive, and sometimes we regret it.

Still it is interesting to reflect on the general question. We collect here some examples. In particular we shall be interested in asking whether or not a given mathematical statement or concept or proof has a natural finite version at all. Sometimes this is not completely clear, and may suggest something new.

Another interesting point concerns the “complexity” of proofs. In classical predicate logic one knows that the size of a proof may have to be very large compared to the size of the statement. In the context of propositional logic this matter is connected to the problem “ $\text{NP} = \text{co-NP}$?” in complexity theory [CR]. (The survey [CS1] provides additional information and references.) One might wonder about the need for infinite processes in a proof which are substantially more complicated than the infinite processes required for the statement. In the context of arithmetic one sometimes measures the infinite complexity of a proof in terms of the strength of the induction that is used. One can also see the matter at play in analysis, geometry, and topology.

Another point about infinite processes is that they lead to more “symmetry”. This symmetry can take a directly geometric form, as in the discussion of the proof of the John-Nirenberg theorem and its finite versions in [CS1], or it can be a symmetry in language. One might ask only whether some quan-

tities are finite or infinite, for instance, and not worry about explicit bounds. This simplification in the language can make it easier to use and formulate “lemmas”, and to find short proofs of theorems through the repeated application of lemmas. This is part of the matter of *cuts* and *cut elimination* in logic, as described in [CS1].

These considerations of logic, complexity, and the lengths of proofs provided part of the motivation for looking more carefully at infinite processes and infinite complexity, but there were other motivations coming from ordinary mathematical activity. It turns out that infinite levels of complexity arise naturally and in some sense unavoidably in geometry and topology, for instance. We shall see more about this later. An enormous amount of classical analysis is about controlling infinite levels of complexity as well.

Of course there are many well-known examples from number theory and combinatorics related to infinite complexity, but we shall emphasize geometry, topology, and analysis here.

It is also pleasant to look for concrete examples from ordinary mathematics to illustrate general phenomena and ideas from formal logic. See [CS2] for further discussions of this nature, with connections with logic more closely drawn.

See [CS3, CS2, CS4] for more discussion about the relationship between short proofs of large objects and the presence of symmetry. We shall return to some of these themes in Section 6.5.

These questions about infinite processes and infinite complexity provide a common thread by which to take a look at mathematics in a broad way. By the end we shall say a little about many different topics, rather than the other way around.

Chapter 2

Examples from the infinite world

Let us look at some concrete examples from ordinary mathematics where infinite processes are involved, perhaps unavoidably.

2.1 Continuity

Consider the notion of continuity, for real-valued functions on the unit interval $[0, 1]$, say. A function $f : [0, 1] \rightarrow \mathbf{R}$ is *continuous* if for every $\epsilon > 0$ and every $x \in [0, 1]$ there exists $\delta > 0$ so that

$$(2.1) \quad |f(x) - f(y)| < \epsilon \quad \text{whenever} \quad |x - y| < \delta.$$

This notion seems to be unavoidably an infinite one. It makes no sense in a finite approximation.

This changes if one prescribes a modulus of continuity, which is to say a particular rate at which the effects of continuity occur. One might consider the Lipschitz condition for a function $f : [0, 1] \rightarrow \mathbf{R}$, which asks that there be a constant $C > 0$ such that

$$(2.2) \quad |f(x) - f(y)| \leq C |x - y| \quad \text{for all } x, y \in [0, 1].$$

One might instead choose an $\alpha \in (0, 1)$ and ask for the weaker condition that there exist a constant C such that

$$(2.3) \quad |f(x) - f(y)| \leq C |x - y|^\alpha \quad \text{for all } x, y \in [0, 1].$$

These provide different ways to measure the continuity of a function. They are not exhaustive; one can give examples of continuous functions which do not satisfy any of these conditions, or any like them.

If we do not specify a rate then it is difficult to see how we can think of continuity intelligently in a finite context. If we choose a particular rate then we can do it. Fix a large integer N , and let us think of the set $I_N = \{\frac{j}{N} : j = 0, \dots, N\}$ as being a finite approximation to $I = [0, 1]$. Given a function $f : I_N \rightarrow \mathbf{R}$, we can look at the smallest constant C for which the analogue of (2.2) is satisfied. This is called the *Lipschitz norm* of f . Alternatively we can fix an exponent $\alpha \in (0, 1)$ and look at the smallest constant C for which the analogue of (2.3) is satisfied. We could fix other rates and measure the approximate continuity for them. As $N \rightarrow \infty$ these measurements can become dramatically different.

For each fixed N no matter what rate we use we are always working with the same functions on I_N , namely all functions. When we pass to the “infinite” realm of functions on I these classes become different. For instance the function $f(x) = \sqrt{x}$ satisfies (2.3) when $\alpha \leq \frac{1}{2}$ but not when $\alpha > \frac{1}{2}$.

There are amusing points here about quantifiers. In the infinite context of the actual unit interval $[0, 1]$ it makes sense to talk about the general notion of continuity, to allow all rates of continuity at once. This universality is accommodated through the use of *universal* and *existential quantifiers*, in the sense of formal logic, and does not work so well for an individual I_N .

In general the infinite context seems to provide a setting where one can work with quantifiers more efficiently. This point interacts with the idea of *cut elimination* from formal logic in any interesting way. (See [CS1] for an introduction.) One knows that quantifiers and their nesting can be conducive to large expansion in cut elimination. One might say instead that the use of quantifiers can help one to make short proofs of complicated statements.

2.2 Uniform continuity

Remember that continuous functions on the unit interval are always uniformly continuous. That is, continuity implies that for each $\epsilon > 0$ there is a $\delta > 0$ so that

$$(2.4) \quad |f(x) - f(y)| < \epsilon \quad \text{whenever} \quad |x - y| < \delta,$$

i.e., there is always a uniform rate at which continuity takes effect. Does this theorem have a natural “finite” version? It would seem not. This is reasonable, since we need to be in the infinite context to make sense of either the assumption or the conclusion. In other words this is a theorem about the relationship between two infinite concepts, with perhaps nothing to say about finite mathematics.

This is a bit troubling.

If we fix a particular (uniform) rate of continuity, as in the Lipschitz condition (2.2), or the Hölder condition (2.3), then we do not really lose information in passing between the infinite context and its finite approximations. For instance, given a Lipschitz function on I , its restriction to each I_N is Lipschitz with the same constant; for large N this restriction determines the original function up to a small error that one can easily compute; any function on I_N can be extended to give a function on I with the same Lipschitz constant, etc. However, having a uniform rate of continuity to begin with is very different from the ordinary notion, which allows different rates at different points.

This leads to another interesting point about continuity: How does one check it in practice? Not in a circular situation, like when one assumes the continuity of two functions and wants to prove the continuity of the sum, but more seriously where an actual function is under consideration. I do not know of a concrete construction which implies the continuity of a function without the possibility to obtain information about the actual rate of continuity that one can somehow write down, at least in principle (and perhaps not easily). The word “concrete” might necessarily exclude many constructions from functional analysis, such as the Hahn-Banach theorem and compactness arguments, as well as transversality arguments and the Baire category theorem. We shall return to these methods later in the text.

In my experience even some not-so-concrete situations lead to particular degrees of continuity. I am thinking for instance of solutions of partial differential equations which are obtained by minimizing some kind of “energy” functional. This means that the function may have been obtained nonconstructively through a compactness argument. Still one can often prove that such a function is continuous, or that its first derivatives are continuous, at least on the complement of a small “singular” set. The methods used in the field that I know do not give continuity with no other information, they tend to provide a condition like (2.3) (at least locally) or they do not work at all. See [Gia], for instance. In many important cases continuity implies

smoothness, and the issue degenerates in the end even if the idea makes sense at an earlier stage of the argument. This does not always happen though, particularly in the more complicated situation where one is trying to study the structure (like smoothness) of a singular set.

Another interesting aspect of continuity is provided by the theorem which says that a continuous function on a compact space always assumes a maximum and a minimum. What is the “finite” version of this? The fact that a function on a finite set always assumes a maximum and a minimum is not very interesting. In some sense continuity and compactness exactly provide the tools to reduce the “infinite” problem to a finite one, and the finite and infinite versions are not really so different.

Alternatively one might think just about the fact that a continuous function on a compact space is automatically bounded, or that if it is positive everywhere then it is actually bounded away from 0. In other words, the existence of uniform bounds which are trivial for finite sets but not in infinite contexts.

Note well the difference between “uniform” and “universal” bounds. A real-valued function on a finite set is always uniformly bounded, but there is no universal bound. The uniform boundedness of continuous functions on compact sets straddles the two.

Sometimes one derives universal bounds in concrete situations through a compactness argument of this type, with no clue as to explicit constants. In practice these arguments typically proceed by contradiction. One assumes that the existence of a sequence for which there is no bound, one uses compactness to extract a subsequence which converges to something, one uses information in the problem to say something about the limit. Effectively one is proving some kind of continuity, but sometimes one can get away with a bit less than that. Arguments of this type tend to be a bit mysterious.

2.3 Classical Banach spaces

A *Banach space* is a normed vector space (over the real or complex numbers) which is *complete* as a metric space (using the metric induced by the norm). Remember that a *norm* on a vector space V is a nonnegative real-valued function $\|\cdot\|$ on V which vanishes only at the zero vector and satisfies the property that satisfies

$$(2.5) \quad \|\lambda v\| = |\lambda| \|v\|$$

for all vectors v and scalars λ , and also the *triangle inequality*

$$(2.6) \quad \|v + w\| \leq \|v\| + \|w\|$$

for all vectors v and w in V . The *metric induced by the norm* is defined by $d(v, w) = \|v - w\|$.

Basic examples include the ℓ^p spaces, $1 \leq p \leq \infty$, which are the spaces of sequences $x = \{x_j\}_{j=1}^\infty$ such that

$$(2.7) \quad \|x\|_p = \left(\sum_j |x_j|^p \right)^{\frac{1}{p}} < \infty.$$

When $p = \infty$ we take ℓ^∞ to be the space of bounded sequences, with norm

$$(2.8) \quad \|x\|_\infty = \sup_j |x_j|.$$

It is well known that these are all norms and that the associated metric spaces are complete.

For $p = \infty$ there is another natural space, namely c_0 , which is the space of sequences $x = \{x_j\}_{j=1}^\infty$ such that

$$(2.9) \quad \lim_{j \rightarrow \infty} x_j = 0.$$

For this space we use the norm $\|\cdot\|_\infty$ again, and one can still show completeness.

Notice that the space of sequences which have zero for all but finitely many coordinates is dense in ℓ^p for $1 \leq p < \infty$. This is not true when $p = \infty$. The closure in ℓ^∞ of this space is precisely c_0 .

Given a Banach space $(X, \|\cdot\|)$ we define its *dual* $(X^*, \|\cdot\|_*)$ to be the space of bounded linear functionals on X . A *linear functional* simply means a linear mapping to the ground field ($\mathbf{R}$ or $\mathbf{C}$ in this case). A linear functional λ on X is said to be *bounded* if there is a constant $M < \infty$ such that

$$(2.10) \quad |\lambda(x)| \leq M \|x\| \quad \text{for all } x \in X.$$

The smallest such M can be written as

$$(2.11) \quad \|\lambda\|_* := \sup_{\substack{x \in X \\ \|x\|=1}} |\lambda(x)|$$

and defines a norm on the space of bounded linear functionals. With this norm $(X^*, \|\cdot\|_*)$ becomes a Banach space in its own right (i.e., it is complete).

If $1 \leq p < \infty$ then the dual of ℓ^p is ℓ^q , where q is determined by the equation $\frac{1}{p} + \frac{1}{q} = 1$. (Thus $q = \infty$ when $p = 1$.) In particular if $1 < p < \infty$ then the dual of the dual of ℓ^p is ℓ^p again. This breaks down for $p = 1, \infty$. The dual of ℓ^∞ is a mess and it is better not to talk about it. However the dual of c_0 is ℓ^1 . Thus the second dual of c_0 is ℓ^∞ , a nice feature which is rather basic in analysis.

What about *finite-dimensional* spaces? For them one can show that the second dual of a space is always the original space again.

Fix an integer n , and define ℓ_n^p in the same way as above, except that we restrict ourselves now to sequences of length n . Thus all of our spaces are just $\mathbf{R}^n$ or $\mathbf{C}^n$ as vector spaces, the differences are in the norms, which we define exactly as above. We cannot define c_0 now, or rather we cannot make a distinction between c_0 and ℓ^∞ ; there is no limit to take. This fits well with the story of duality, because each of ℓ_n^1 and ℓ_n^∞ is the Banach-space dual of the other. The content in this statement lies in the norms, since the underlying vector spaces and their duals are just $\mathbf{R}^n$ or $\mathbf{C}^n$.

Thus again we have some distinctions in the “infinite” case which do not make sense in the “finite” approximations.

Although in finite dimensions the second dual of a space always gives the space back, one can make quantitative distinctions between the cases of $p = 1, \infty$ and $1 < p < \infty$ for the ℓ_n^p 's. For instance, the norms for $1 < p < \infty$ enjoy stronger convexity properties which persist to infinite dimensions.

See [HSt, Ru2, Ru3] for more information about these topics (at an introductory level).

2.4 The open mapping theorem

Theorem 2.11 *Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be Banach spaces, and suppose that $T : X \rightarrow Y$ is a bounded linear mapping. (Boundedness means that there is a constant $M > 0$ so that*

$$(2.12) \quad \|T(x)\|_Y \leq M \|x\|_X \quad \text{for all } x \in X,$$

and this condition is equivalent to continuity in the case of linear mappings.) If the image of T is all of Y , then T is an open mapping, i.e., it sends open subsets of X to open subsets of Y .

This is a well-known theorem due to Banach. (See [Ru2].) The main point about the assumption of surjectivity is that the image of T be closed. If it is closed then one can always treat the image as a Banach space in its own right.

To understand the meaning of this theorem it is helpful to consider some special situations. The first point is that the theorem does not have much content for finite-dimensional spaces. In finite dimensions all norms give equivalent topologies, and the theorem reduces to a simple observation about linear mappings between Euclidean spaces. In infinite dimensions it is much more substantial.

What does the condition that T be open really mean? Using linearity one can show that T is an open mapping if and only if $T(B_X(0, 1))$ contains the origin of Y in its interior, where $B_X(0, 1)$ denotes the unit ball in X . In other words, one only has to test for openness “once”. This is the same as asking that there exist an $\eta > 0$ so that

$$(2.13) \quad T(B_X(0, 1)) \supseteq B_Y(0, \eta).$$

Assume now that T is one-to-one. In this case (2.13) is equivalent to the lower bound

$$(2.14) \quad \|T(x)\|_Y \geq \eta \|x\|_X \quad \text{for all } x \in X.$$

This is very nice. The lower bound (2.14) implies that the kernel of T is trivial, and it is a kind of quantitative version of injectivity. It is an automatic consequence of injectivity in finite dimensions but not in infinite dimensions, and the content of the open mapping theorem is that in infinite dimensions (2.14) follows from injectivity if one also knows surjectivity.

As noted above surjectivity can be replaced by the requirement that the range of T be closed, and indeed (2.14) forces the range to be closed. The latter assertion is a standard exercise; the (metric) completeness of the domain implies the completeness of the image, under the assumption of a lower bound like (2.14). Thus one can also think of (2.14) as being a quantitative version of the requirement that the range be closed.

The open mapping theorem can therefore be viewed as a criterion for the existence of a uniform bound. As such it is a bit like the fact that continuous functions on compact spaces are always uniformly bounded.

To make the role of the uniform bound more concrete let us consider a special case. Take $X = Y = \ell^p$ as defined in the previous section, with any (single) choice of $1 \leq p \leq \infty$. We could also take $X = Y = c_0$ instead.

Fix a sequence of scalars $\{t_j\}_{j=1}^\infty$ which are bounded and never 0. Define an operator T by

$$(2.15) \quad T(x) = y, \quad y_j = t_j x_j \quad \text{for all } j.$$

That is, x and y are sequences, and T is a “diagonal” operator with entries t_j . This defines a bounded linear operator on $X = Y = (\ell^p \text{ or } c_0)$. The requirement that the t_j ’s never vanish is the same as the injectivity of the operator T , and it is also equivalent to the requirement that the range of T be dense in this special case of diagonal operators. One can check that (2.14) is the same as

$$(2.16) \quad |t_j| \geq \eta \quad \text{for all } j$$

in this case. Thus the open mapping theorem provides a criterion for a uniform lower bound on the size of these scalars t_j .

If we were to make the analogous construction in finite dimensions, using scalars $\{t_j\}_{j=1}^n$ and defining T on ℓ_n^p , we would simply say that (2.16) holds as soon as we know that the t_j ’s are all different from 0. In finite dimensions surjectivity is the same as having dense image for linear mappings. In infinite dimensions these properties are more subtle and involve infinite processes. The effect of the open mapping theorem is to relate two different kinds of infinite processes in a nontrivial way.

Is the open mapping theorem useful? It is frequently very important in analysis to be able to derive lower bounds like (2.14). In my experience however the open mapping theorem has been primarily of psychological value. The main approach that an analyst has to proving something like the surjectivity of a mapping is to first establish an inequality like (2.14). The open mapping theorem implies in some sense that there is no other way to do it.

2.5 Category versus measure

In mathematics it is very convenient to have reasonable ways to talk about “almost all” points in a space. There are two especially famous ways to do this. The first is to use the Baire category theorem. In a complete metric space we might think of a set E which is a countable intersection of dense open sets as containing almost all elements of the original space. Baire’s theorem implies that any such E is always dense. In particular the intersection of any two or even countably many of these putatively large sets remains dense and in the same class of subsets.

Another possibility is to work on a measure space and to look at sets whose complement has measure zero. Again we can take *countable* intersections and still have a set of the same type.

In some cases one can use algebraic or analytic structure to make much more refined notions of “almost all” points in a space. One can use proper subvarieties to make notions of negligibility for subsets.

A more primitive idea is to use cardinality, to simply look at subsets whose complement has cardinality bounded in a certain way. Inside an infinite set one might consider subsets whose complement is finite. For uncountable sets one can look at subsets whose complement is at most countable.

In metric spaces one can impose more precise “smallness” conditions on the complement of a set, through Hausdorff measure or Minkowski content, for instance. In topology one might use submanifolds of smaller dimension, or just subsets of smaller topological dimension. (These concepts will be defined later in the text.)

The idea of “almost every” cannot work in the finite world in a similarly nice way. We cannot have notions of “almost every” which are closed under arbitrary finite intersections and which are also nontrivial. The empty set should not contain almost all the elements of a nonempty set! What can we do in finite contexts?

One possibility is to replace “zero measure” with “small measure”, and then restrict the kind of intersections that one makes. This is awkward but feasible nonetheless, and commonplace in counting arguments. The Baire category theorem is more problematic. I do not see a natural version for finite mathematics. This is slightly disturbing, particularly for the idea of using Baire category as a method for proving existence of points with special properties.

The proof is less scary. One “finds” actual sequences which converge to a point in the intersection. The method is fairly concrete even if it is not so clear how to make more concrete statements. To analyze it carefully one can think about trying to implement it on a computer, to actually produce the relevant sequence. We then have to implement our “assumptions” that the initial sets are dense and open.

This is not ridiculous, just very awkward. Here again we see how quantifiers can be manipulated much more efficiently in “infinite” contexts, and how much more simple and “symmetric” definitions can be in the infinite world. The notions of openness and density rely on quantifiers very explicitly. One also sees some of the cycling associated to proofs with cuts and

quantifiers.

For the record let us sketch the proof. We have a complete metric space $(M, d(x, y))$ and a sequence $\{U_j\}_{j=1}^\infty$ of dense open sets, and we want to show that $\bigcap_j U_j$ is also dense.

A basic lemma is that $V_j = U_1 \cap U_2 \cap \dots \cap U_j$ is dense and open for each j . One can check that the intersection of a pair of dense open sets is dense and open, and then repeat. Although simple to prove this part is already substantial, in the sense that dense open sets provide a cruder notion of “almost all” points in a metric space with good properties. This cruder notion also lacks an obvious counterpart for finite mathematics, and enjoys a kind of symmetry which is useful in making proofs.

Next one chooses a sequence of points $\{x_j\}_{j=1}^\infty$ in M and a sequence of positive numbers $\{\epsilon_j\}_{j=1}^\infty$ with the following properties:

1. each x_j lies in V_j , and in fact $\overline{B}(x_j, 2\epsilon_j) \subseteq V_j$,
2. $d(x_{j+1}, x_j) \leq \epsilon_j$ for all j , and
3. $\epsilon_{j+1} \leq \frac{\epsilon_j}{2}$ for all j .

It is not difficult to make these choices, using the fact that $\{V_j\}_{j=1}^\infty$ is a decreasing sequence of dense open subsets. They imply that

$$d(x_j, x_k) \leq 2\epsilon_j \quad \text{when } k \geq j,$$

by summing a geometric series. Thus $\{x_j\}_{j=1}^\infty$ is a Cauchy sequence in M , and therefore converges, and the properties above ensure that the limit lies in all the V_j 's.

Note that the notions of “almost every” provided by Baire category and measure need not be compatible with each other, even on the real line using Lebesgue measure. This is true in principle and in actual practice.

If one wanted to analyze Baire category arguments more carefully, looking for more quantitative versions, or just to bring out some kind of constructivity, then a natural class of spaces to consider would be the following. Let $\{F_j\}$ be some sequence of *finite* sets, and let $\mathcal{F}$ denote their Cartesian product. Thus an element $x \in \mathcal{F}$ actually represents a sequence $\{x_j\}$, for which $x_j \in F_j$ for each j . This space can be given a topology in the usual way, where two elements x and y are “close” if their entries $x_j, y_j \in F_j$ agree for a large number of j 's. This is the usual *product topology*, and in this case it

defines a space which is compact and for which the topology is determined by a metric.

Now in this case the properties of being dense or open admit more concrete interpretations than usual, and one can analyze the argument above to make it more concrete.

Chapter 3

Symmetry and language

There is a common thread to the examples above, concerning “abstract” concepts in infinite contexts and the possibility for meaningful “concrete” versions in finite situations. Here is a sampling.

- We have *continuity* and *uniform continuity* for functions, and then more concrete versions where a *rate* of continuity is specified, as in the Lipschitz and Hölder conditions.
- We can talk about the *boundedness* of linear mappings between Banach spaces, and then more concretely we can look at the *size* of the bounds. In particular *all* linear operators between finite-dimensional spaces are bounded.
- We can think of a subset of a space as being negligible if it has measure zero, while in the finite world we might have to be more careful and consider the actual number of elements of the subset as compared to the ambient space as a whole.
- In topological spaces we can look at the properties of being *dense* and *open*, for which elegant concrete versions are not apparent.
- We can consider merely the *finiteness* or *infiniteness* of some quantity, as opposed to trying to decide which numbers are *large* and which are *small*.

This list is by no means complete, but it gives a flavor. The extent to which the infinite concepts can be captured by finite versions depends on the details of the situation.

The extent of abstractness or concreteness used has important consequences for the enterprise of making proofs. In mathematics we consider something to be “known” when we have proved it and not otherwise. What if something is true but all proofs are very long?

Mathematicians have long been concerned about matters of undecidability. We have the goal of “truth”, we try to formalize it, we make axiom systems to cover what we are interested in, and then some questions are typically left over. This is not too surprising after all, but we often enter the field as students with more idealism and naïveté about the notion of “truth”. There are even precise theorems of Gödel and others to the effect that this is unavoidable in the context of arithmetic. (See [Man], for instance.) There are even quantitative versions of Gödel’s incompleteness theorems which have been given by Chaitin, based on algorithmic information theory.

In simpler forms of logic we do not have this problem. Propositional and first-order predicate logic are consistent and complete. “All is well”, in a sense.

But is it really? What about “practical” incompleteness, where a statement is true but all the proofs are too enormous for a human being to find?

In predicate logic it is well known that the set of tautologies form an *algorithmically undecidable set*. This means that there is no computer program which can input a sentence in (first-order) predicate logic and decide whether or not the sentence is a tautology. There are computer programs that can generate a complete list of all (first-order) tautologies; one can simply list all strings that could possibly represent a proof, check which ones do represent a proof, keep those and throw away the rest. The problem is to know whether a given sentence will appear on the list eventually or not, and for that there is no such algorithm. See [EFT], for instance.

As a practical matter this means that there can be tautologies whose proofs are enormously larger in size than the size of the statement itself. “Enormously” means here larger than any recursive function in the size of the statement of the tautology.

There are similar issues for propositional logic (i.e., classical logic but without quantifiers, relations, etc.). In that case the set of tautologies is decidable and quite easily so, one has only to check truth tables. The problem is that the naïve algorithm could take an exponential amount of time (in the size of the given propositional formula) to decide whether the formula is a tautology or not. Exponential expansion is mild compared to nonrecursive growth, but still prohibitive practically.

There is an algorithm which can determine in polynomial time whether a given propositional formula is a tautology if and only if $P = NP$. The latter is a famous question in complexity theory; see [GJ, HU, Pap] for background information about it and related problems of algorithmic complexity. The more precise statement is that the set of propositional formulas which are *not* tautologies form an NP-complete set. As mentioned in Chapter 1, there is a proof system in which propositional tautologies always have polynomial-sized proofs if and only if $NP = co-NP$; see [CR] for more details.

Even in propositional logic one has therefore serious difficulty with being able to find proofs of modest size for all tautologies.

What is it that allows one to make short proofs? In [CS3, CS2, CS4] there is an idea put forward to the effect that the existence of short proofs of complicated statements reflects the presence of some kind of underlying symmetry. This might be a literal “geometric” symmetry of some object, but it might also appear as a symmetry in the language, in the way that objects are described. In our exploration of infinite processes and infinite concepts in mathematics we see many examples which illustrate this idea from [CS3, CS2, CS4].

It is easier to make reasoning about the “abstract” concept of finiteness or infiniteness of some quantity than it is to make more precise estimates. It is easier to formulate lemmas based on the distinction between finiteness and infiniteness, and to use the same lemma many times.

These are familiar themes to any working mathematician. One knows from practical experience that it is easier to work with the notion of *continuity* of functions than it is to keep track of how the modulus of continuity changes under various operations, and that it is easier to say that a set is small because it is the countable union of sets of measure zero than to make some concrete measurement of its negligibility.

Thus it can be convenient to work with infinite processes or concepts even if in the end we are really interested in something more concrete. The relevant distinctions (between large and small, for instance) can simplify in the infinite world; we might ask for the mere *existence* of a bound, without trying to keep track of what it is. This makes it easier to formulate and use lemmas and then to find actual proofs.

After we find a proof we can try to go back and analyze it to get explicit bounds, or at least to say that such bounds are implicit in the argument. This idea is made general and precise in formal logic through *cut elimination*, which in effect seeks to transform a given proof into a “direct” one, with no

lemmas, each step being explicit for the particular case at hand. See [CS1] for an introduction and [Gir] for more details.

Chapter 4

The existence of limits

There are many results in classical analysis and related areas of mathematics which assert the existence of a limit, perhaps only “almost everywhere”. To what extent do these statements have natural finite versions? Does their content lie purely within the infinite world?

4.1 The Riemann-Lebesgue lemma

Let $f(x)$ be an integrable function on the real line. Define its *Fourier transform* $\hat{f}$ by

$$(4.1) \quad \hat{f}(\xi) = \int_{\mathbf{R}} f(x) e^{-2\pi i x \xi} dx.$$

Roughly speaking, the Fourier transform tells us how to decompose a function into primitive “frequencies”. This idea is made precise by the “inversion” formula

$$(4.2) \quad f(x) = \int_{\mathbf{R}} \hat{f}(\xi) e^{2\pi i x \xi} d\xi.$$

This formula works when $\hat{f}$ is also integrable, but in general it requires some interpretation. See [HSt, Ru2, Ru3, SW] for more information.

The following is a basic fact in Fourier analysis.

Theorem 4.2 *For each integrable function f on $\mathbf{R}$ we have that*

$$(4.3) \quad \lim_{|\xi| \rightarrow \infty} \hat{f}(\xi) = 0.$$

An annoying feature of this theorem is that we cannot say anything about how fast the Fourier transform decays at infinity. And indeed we *cannot* say anything, because the rate of decay can be arbitrarily slow. What really is the content of the theorem then?

The first basic observation is that the Fourier transform is uniformly bounded under these conditions. Namely,

$$(4.4) \quad \sup_{\xi \in \mathbf{R}} |\hat{f}(\xi)| \leq \int_{\mathbf{R}} |f(x)| dx.$$

This fact is very robust in terms of making finite approximations.

The second observation is that for “nice” functions one can get decay that is easily measured. There is a general principle to the effect that smoothness of a function is reflected in decay at infinity of the Fourier transform, and there are many specific results of this nature (often proved by integrating by parts). Let us content ourselves with a special case which is easy to calculate. Let $a, b \in \mathbf{R}$ be given, with $a < b$, and consider the function

$$(4.5) \quad g_{a,b}(x) = \frac{1}{b-a} \mathbf{1}_{[a,b]}(x).$$

Here $\mathbf{1}_A(x)$ denotes the characteristic (or indicator) function of the set A , which means the function which takes the value 1 on the set and 0 off the set. The normalizing factor of $\frac{1}{b-a}$ was included so that

$$(4.6) \quad \int_{\mathbf{R}} |g_{a,b}(x)| dx = 1.$$

The Fourier transform of this function is easy to compute. It is given by

$$(4.7) \quad \hat{g}_{a,b}(\xi) = \frac{1}{b-a} \frac{1}{-2\pi i \xi} (e^{-2\pi i b \xi} - e^{-2\pi i a \xi}).$$

Notice that this is compatible with (4.4) even if it may not appear to be at first. The potential for singularities provided by the denominators is avoided by the difference in the exponentials.

For all a, b, ξ we have that

$$(4.8) \quad |\hat{g}_{a,b}(\xi)| \leq \frac{2}{b-a} \frac{1}{2\pi|\xi|}.$$

That is, we have bounded the difference the exponentials by 2. For “most” ξ ’s the difference in the exponentials will not be small. It will be small only when $(b - a)\xi$ is close to an integer.

This inequality makes concrete the second observation. If f is actually a step function – a finite union of characteristic functions of intervals – then the Fourier transform decays at infinity like a multiple of $\frac{1}{|\xi|}$.

To prove the theorem one combines these two pieces of information using a density argument, as follows. Suppose that we are given an integrable function $f(x)$ on $\mathbf{R}$, and let $\epsilon > 0$ be given. A standard fact from real analysis is that there is a step function g on $\mathbf{R}$ such that

$$(4.9) \quad \int_{\mathbf{R}} |f(x) - g(x)| dx < \epsilon.$$

Using (4.4) we get that

$$(4.10) \quad \sup_{\xi \in \mathbf{R}} |\hat{f}(\xi) - \hat{g}(\xi)| < \epsilon.$$

On the other hand our earlier calculations show that

$$(4.11) \quad \lim_{|\xi| \rightarrow \infty} \hat{g}(\xi) = 0.$$

This implies that

$$(4.12) \quad \limsup_{|\xi| \rightarrow \infty} |\hat{f}(\xi)| \leq \epsilon.$$

Since $\epsilon > 0$ is arbitrary we conclude that the limit actually vanishes.

We cannot say anything about the rate at which the Fourier transform decays at infinity. There are two reasons for this. Although we can approximate an integrable function f by a step function g as above, we cannot say much about the nature of the approximation, i.e., how will the complexity of g grow as ϵ shrinks to 0? Without additional assumptions on f the answer is that it can grow as fast as it wants.

The second reason is that in our example above we had the factor of $\frac{1}{b-a}$ in (4.8). This can be made as large as we want by taking b close to a , without affecting the L^1 norm of $g_{a,b}$. We saw also that for most ξ we cannot really do much better than the estimate in (4.8). Of course we can build much nastier integrable functions by combining infinitely many functions like $g_{a,b}$ with $b - a$ as small as we want. We have only to be careful to include a summable series of coefficients to get an integrable function in the end.

The bottom line is that the rate of decay of the Fourier transform of a general integrable function is unknowable because it depends on the structure of a general integrable function, which can be practically arbitrary. In effect there are infinite processes in both the statement and the conclusion of the theorem.

We also saw another important point: that a crucial ingredient for the result about the limit was the uniform bound (4.4), which does have a more “absolute” meaning. In particular it works perfectly well in finite approximations.

4.2 Lebesgue's theorem

Suppose that f is an integrable function on $\mathbf{R}^n$. Then

$$(4.13) \quad \lim_{r \rightarrow 0} \frac{1}{|B(x, r)|} \int_{B(x, r)} f(y) dy = f(x)$$

for almost all $x \in \mathbf{R}^n$. Here $|B(x, r)|$ denotes the volume of the ball $B(x, r)$. Thus we are taking the average of the values of f on the ball of radius r around x , then shrinking the radius to 0, and we are looking for the limit to be the original value at that point. This would occur at all points if f were continuous, but for functions which are merely integrable the limit need not even exist at a given point.

What can we really say about this limiting process? What does it mean concretely? Is there any “finite” information contained in this result?

As in the previous section one cannot say anything about the rate at which the limit converges. One could not say anything even for continuous functions. This ambiguity is built into the statement, because the continuity or integrability of a function does not give terribly precise information about its structure.

How can one prove such a theorem? The basic structure of the argument is much the same as before. A key point is the density of continuous functions. Given an integrable function f on $\mathbf{R}^n$ and an $\epsilon > 0$ there is a *continuous* function g on $\mathbf{R}^n$ such that

$$(4.14) \quad \int_{\mathbf{R}^n} |f(x) - g(x)| dx < \epsilon.$$

We want to use this to show that the existence of limits for continuous functions implies the corresponding result for integrable functions.

To do this we need to have a uniform bound, analogous to (4.4) in the previous situation. In this case the geometry is more tricky. Consider the *maximal function*

$$(4.15) \quad f^*(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy.$$

In general when one is interested in the existence of a limit it is helpful to first obtain uniform bounds. This case is more complicated because we need to take the supremum over r for each point x , but we need to treat the points x more individually (and not just take the supremum).

Suppose for instance that we knew that there was a constant $C > 0$ so that

$$(4.16) \quad \int_{\mathbf{R}^n} f^*(x) dx \leq C \int_{\mathbf{R}^n} f(x) dx$$

for all integrable functions f on $\mathbf{R}^n$. Then we would be able to show that (4.13) holds almost everywhere for integrable functions using the fact that it is true for continuous functions and an approximation argument, in much the same manner as in the previous section.

Unfortunately (4.16) is wrong. In fact, as soon as f is not 0 almost everywhere, one has that

$$(4.17) \quad f^*(x) \geq \frac{a}{1+|x|}$$

for some constant $a > 0$ (which depends on f). This is not hard to verify, and it implies that

$$(4.18) \quad \int_{\mathbf{R}^n} f^*(x) dx = \infty.$$

The analogue of (4.16) for L^p spaces is true for all $p > 1$, though. There is a substitute for $p = 1$, which says that f^* is not too big most of the time. The precise statement is that there is a constant $C > 0$ such that

$$(4.19) \quad \lambda \cdot |\{x \in \mathbf{R}^n : f^*(x) > \lambda\}| \leq C \int_{\mathbf{R}^n} f(x) dx$$

for all integrable functions f on $\mathbf{R}^n$. Note that (4.19) would follow automatically if (4.16) were true.

It turns out that this weaker version is sufficient for proving that (4.13) holds almost everywhere for integrable functions. See [Ga, St2, SW, Se9].

There are several nice points now. The first is that an estimate like (4.19) makes perfectly good sense in discrete approximations. In fact (4.19)

is equivalent to having uniform bounds for its analogues in finite approximations. The theorem about the existence of limits does not directly make sense in a finite context, but these bounds do make sense, and they are a large part of the story.

Actually there are ways to bring the existence of the limit back to the finite world. Although one cannot control the rate at which the limit converges, one can count the number of oscillations. That is, one can fix an $\epsilon > 0$ and ask how many times the average

$$(4.20) \quad \frac{1}{|B(x, r)|} \int_{B(x, r)} f(y) dy$$

oscillates by more than ϵ as r shrinks from 1 down to 0, say. If the limit as $r \rightarrow 0$ exists then the number of these oscillations must be finite. One can try to derive bounds for this number, as a function of x . Bounds on averages of these numbers, or on the measure of the set of x 's where they are large, as in (4.19). Such results can be derived using the methods of Carleson's Corona construction, and results like this are explained (in some versions) in [Ga]. Again these bounds are equivalent to the existence of uniform bounds in finite contexts.

Thus in fact there are "finite" statements which correspond to existence almost everywhere of limits, even if there are problems with the most obvious ways to do this.

Another interesting aspect is that in many situations a maximal estimate like (4.19) is necessary to have existence of limits almost everywhere. That is, there are actual theorems to the effect that existence of limits imply suitable bounds. See [St1, N, Mau]. This fits nicely with the question of what really exists at the level of "finite" mathematics. It is typically easier to make sense of the idea of a bound in finite contexts than the existence of the limit, and although the two appear to be quite different, in fact they are closely connected.

There is a simpler and older result of this type in the context of linear mappings between Banach spaces, called the uniform boundedness principle. (See [Ru2].) Suppose that you have a sequence of bounded linear mappings T_j from some Banach space $(X, \|\cdot\|_X)$ into another one $(Y, \|\cdot\|_Y)$. Suppose also that

$$(4.21) \quad \sup_j \|T_j(x)\|_Y < \infty \quad \text{for all } x \in X.$$

Then the T_j 's have uniformly bounded operator norms, which means that

there is a constant $M < \infty$ such that

$$(4.22) \quad \sup_j \|T_j(x)\|_Y \leq M \|x\|_X \quad \text{for all } x \in X.$$

Thus pointwise bounds imply uniform bounds, which is slightly remarkable. The point here is that X is allowed to be infinite-dimensional. The theorem would be pretty trivial otherwise.

Note that (4.21) holds automatically when

$$(4.23) \quad \lim_{j \rightarrow \infty} T_j(x) \text{ exists for all } x \in X.$$

In practice it is not so easy to prove (4.23) without first establishing a uniform bound.

The uniform boundedness principle provides another example of a theorem which in effect seems to relate two kinds of infinite processes in a nontrivial way while not appearing to have a natural finite version.

4.3 The ergodic theorem

There is a variation on the themes of the preceding section which are closer to physical reality. Let $(X, \mathcal{M}, \mu)$ be a finite measure space, and let $T : X \rightarrow X$ be a measurable mapping which is measure-preserving, so that

$$(4.24) \quad \mu(T(A)) = \mu(A) \quad \text{for all measurable sets } A \subseteq X.$$

For example, think of X as being the unit circle equipped with ordinary Lebesgue measure, and let T be rotation by the angle θ .

Fix a point $x \in X$, and consider the sequence $\{T^j(x)\}_{j=1}^{\infty}$ of images of x under T . This is called the *orbit* of x under T . How should we expect it to be distributed in X ?

This question arises in classical mechanics. Think of a particle in a physical system moving under the influence of some conservative force field. For this one should think in terms of continuous time instead of discrete time, as we have here, but this is not a serious issue. For instance, one can fix a small number $\delta > 0$, and take T to be the mapping that tells how a particle described by x moves after flowing in the system by an amount of time equal to δ .

In the context of mechanics the assumption that T be measure-preserving is very natural, for reasons that we shall not discuss. (See [Ar].) For this we should say that the phase space X tracks not only the position of the particle but also its momentum.

One might hope that the orbit of a point is uniformly distributed throughout the space. This is not true in general, and one can make easy counterexamples. It is true however for *almost all* orbits when T is an *ergodic* transformation. This assumption says that if $A \subseteq X$ is measurable, and if $T(A) = A$, then either $\mu(A) = 0$ or $\mu(A) = \mu(X)$. A subset A such that $T(A) = A$ is said to be *invariant*. The existence of nontrivial invariant subsets would prevent the orbits from being uniformly distributed, and so this assumption is at least necessary.

To understand what this means it is helpful to come back to our example of a rotation on the unit circle by an angle θ . If θ is a rational multiple of π then the corresponding rotation is not ergodic, and this is easy to check. In this case the mapping is even periodic, so that T^n is the identity for some positive integer n , and all orbits are finite. It turns out that the rotation is an ergodic transformation when the angle is an irrational multiple of π . See [Sn] this and other more difficult examples.

The precise formulation of the *ergodic theorem* states that if $T : X \rightarrow X$ is measure-preserving and ergodic, and if f is an integrable function on X , then

$$(4.25) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N f(T^j(x)) = \int_X f d\mu \quad \text{for } \mu\text{-almost all } x \in X.$$

In other words, if we average f over X , that is the same as taking a random point $x \in X$ and averaging f over the orbit of x under T . Sometimes one says “time averages equal space averages”.

For example, we might take f to be the characteristic function of a measurable set A . In this case the left side of (4.25) measures the amount of time that the orbit of x spends in A , while the right side reduces to the measure of A . Note that the existence of the limit on the left side is part of the conclusion.

This theorem is reminiscent of the existence of limits almost everywhere in the preceding section, except that it goes in the opposite direction, making averages in the large instead of over small balls. This turns out not to be a major issue. For instance one of the main ingredients in the proof is a

estimate for the maximal function

$$(4.26) \quad \sup_N \frac{1}{N} \sum_{j=1}^N |f(T^j(x))|$$

which is analogous to the one given before in (4.19).

See [Bi] for a proof of the ergodic theorem and further information. See [CW] for techniques for relating this more closely to the matter of the previous section, through the method of “transference”.

If the mapping is measure-preserving but not ergodic one still has bounds for the maximal function and the existence almost everywhere of the time averages. One simply cannot identify the time averages with the space averages.

As usual the existence of a limit does not have “finite” meaning but much of the mathematics around the theorem does have finite meaning. In this case however the physical interpretation makes the question of finite meaning more compelling and perhaps more intuitive or attractive. For this one should look at more specific situations and their dynamics, and this leads to a large area of current mathematical research.

4.4 Differentiability almost everywhere

Let $f : \mathbf{R}^n \rightarrow \mathbf{R}$. We say that f is differentiable at the point p if there is a linear mapping $A : \mathbf{R}^n \rightarrow \mathbf{R}$ (the differential of f at p) such that

$$(4.27) \quad f(x) = f(p) + A(x - p) + \epsilon(x),$$

where

$$(4.28) \quad \lim_{x \rightarrow p} \frac{\epsilon(x)}{|x - p|} = 0.$$

There is a famous old theorem to the effect that Lipschitz functions are differentiable almost everywhere. (See [Fe, St2, Se9].) As in Section 2.1, a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is said to be *Lipschitz* if there is a constant $C > 0$ so that

$$(4.29) \quad |f(x) - f(y)| \leq C |x - y| \quad \text{for all } x, y \in \mathbf{R}^n.$$

This theorem is quite remarkable. Although it can be seen as a variant of the one discussed in Section 4.2, it is noticeably different. In Section

4.2 we started with an integrable function f and we wanted to show that the averages of the values of f on small balls centered at a random point x converge to the value of f at x . In particular we had a candidate for the limit in our hands already. To prove that Lipschitz functions are differentiable almost everywhere, part of the problem is to find the candidate for the limit.

Another point about this theorem is that Lipschitz functions are arguably more “elementary” than integrable functions. Just the definition of Lebesgue integrals, integrability, and implicitly measurability entail infinite processes of some sophistication. In the Lipschitz condition we have only to take a supremum over all pairs of points and ask that the result be finite. This is also an infinite process, but it is more straightforward.

The existence of a derivative of a Lipschitz function at a point provides a kind of “finiteness” to its behavior. It says that the function is approximately affine at the given point, and affine functions span a finite-dimensional space. The asymptotic behavior of Lipschitz functions on general metric spaces need not have “finite dimensional” limitations on their asymptotic behavior of this type.

Although one has the existence almost everywhere of the derivative of a Lipschitz function, one cannot say anything about the rate at which the convergence takes place. How then might look for natural versions of the theorem in finite contexts?

Normally we might ask for a uniform bound of the objects whose limit we wish to take, but in this case we have simply assumed it in (4.29). (There is a different version of the story in which uniform bounds on the difference quotients are not so immediate and are more crucial to the existence of the limit, but we shall not pursue that here.)

Instead we can look for good bounds on the oscillation of the function. For instance, we can take a ball $B(x, t)$ and ask how well f is approximated by an affine function on this ball, as in

$$(4.30) \quad \alpha(x, t) = \inf_A \sup_{y \in B(x, t)} \frac{|f(y) - A(y)|}{t},$$

where the infimum is taken over all affine functions A . For almost all x we have that

$$(4.31) \quad \lim_{t \rightarrow 0} \alpha(x, t) = 0,$$

because of the differentiability almost everywhere. One can look for quantitative bounds on the $\alpha(x, t)$'s which would make sense in discrete approxi-

mations. Such bounds exist, as in [Do]. These bounds involve averages over x and t , and not uniform rates of convergence.

In these bounds we would permit ourselves to approximate f on balls by affine functions that are turning around as we move the balls, but in fact one can also control the manner in which these affine approximations are turning around. This is similar to the story of Carleson's Corona construction [Ga], as mentioned before in Section 4.2.

These points about uniform bounds are discussed further in [DS3, Se5, Se9].

The bottom line is that there are actually estimates which work for discrete approximations and which capture rigidity properties of Lipschitz mappings.

Chapter 5

Using infinite processes

Many times in mathematics it is very convenient to use infinite processes, ones which are apparently much more complicated than the problems to which they are applied. We give some examples here in this chapter, with emphasis on geometry and topology. We are particularly interested in situations where the statement is already pretty concrete even if the proof is not. The matter is not as much fun when the statements themselves already rely on abstract processes.

These examples should be compared to the discussion of lengths of proofs in Chapter 3.

5.1 Mostow rigidity

We consider a rigidity phenomenon which is closer to differential geometry and Lie groups. For this discussion it will be convenient to have the idea of *hyperbolic geometry*. The easiest way to describe this is to use a basic model, the upper half-space $\mathbf{R}_+^n$, which means the set of points whose last coordinate is positive:

$$(5.1) \quad \mathbf{R}_+^n = \{x \in \mathbf{R}^n : x = (x_1, \dots, x_n), x_n > 0\}.$$

We can define a *Riemannian metric* on this set using

$$(5.2) \quad \frac{dx_1^2 + \dots + dx_n^2}{x_n^2}.$$

One can think of this simply as a recipe for computing infinitesimal lengths of curves. The lengths are distorted by a factor of x_n^{-1} as we move towards

the boundary hyperplane $x_n = 0$. “Distorted” means relative to ordinary Euclidean geometry.

To give a feeling for the behavior of this geometry, consider the transformation

$$(5.3) \quad x \mapsto a x,$$

where a is a positive real number. This transformation has the simple effect on Euclidean geometry of changing distances by exactly a factor of a . As for our hyperbolic metric, notice that transformations of this form map $\mathbf{R}_+^n$ onto itself, and that *they do not change hyperbolic distances*. One can use this together with the obvious invariance of the metric under translations $x \mapsto x + b$, $b \in \mathbf{R}^n$, $b_n = 0$, to show that *all points in $\mathbf{R}_+^n$ look the same for the hyperbolic metric*. That is, given any pair of points in $\mathbf{R}_+^n$, one can find a mapping from $\mathbf{R}_+^n$ onto itself which preserves the hyperbolic geometry.

We shall restrict ourselves to $n \geq 2$ for this discussion. Although the definition makes sense for $n = 1$, it does not give back anything new, just another copy of the Euclidean line. The mapping $x \mapsto e^x$ provides the correspondence between the two metrics.

We can define hyperbolic geometry as a general concept using this model. Let M be a smooth manifold of dimension n . If we have chosen a Riemannian metric on M , then we call this metric *hyperbolic* if around each point in M there is a coordinate chart which identifies a neighborhood of the point in M with a neighborhood of a point in $\mathbf{R}_+^n$ in such a way that the metric on M corresponds exactly to our metric on $\mathbf{R}_+^n$. There are various ways to formulate this, in terms of “constant negative curvature”, for instance, but they amount to this in the end.

An amazing theorem of Mostow [Mos] implies that if M_1 and M_2 are two compact hyperbolic manifolds of the same dimension n whose fundamental groups are isomorphic, and if $n > 2$, then M_1 and M_2 are *isometrically equivalent*. This means that there is a mapping from M_1 onto M_2 which preserves the metric. This is extremely wrong in dimension 2, in which surfaces can admit *continuous* families of inequivalent hyperbolic geometries.

What does this theorem mean in terms of “finite” mathematics?

To understand this question better it is helpful to consider a reformulation. If M is a compact hyperbolic manifold of dimension n , then it can be realized as a quotient of our model space $\mathbf{R}_+^n$. That is, there is a discrete group Γ of isometries of $\mathbf{R}_+^n$ such that M is isometrically equivalent to $\mathbf{R}_+^n/\Gamma$. The *isometries* of $\mathbf{R}_+^n$ are the mappings from $\mathbf{R}_+^n$ onto itself which preserve

the hyperbolic metric. This is a group, and in fact it can be described as a classical Lie group of matrices. It can also be realized as the group generated by translations and dilations mentioned above together with reflections about spheres whose centers lie on the hyperplane $x_n = 0$.

This representation of M as $\mathbf{R}_+^n/\Gamma$ is not terribly mysterious. Another way to say it is that the universal covering of M is isometrically equivalent to $\mathbf{R}_+^n$. By assumption M is locally equivalent to $\mathbf{R}_+^n$, and so the point is to knit together many local equivalences into a global one. By passing to the universal covering we avoid ambiguities that could come from having a sequence of local equivalences loop around to a place where they have been before, but where a different choice of local equivalence was used.

At any rate this representation is well known. The group Γ is isomorphic to the fundamental group of M as a discrete group. Thus in the context of Mostow's theorem we can represent M_1 and M_2 as $\mathbf{R}_+^n/\Gamma_1$ and $\mathbf{R}_+^n/\Gamma_2$, where Γ_1 and Γ_2 are groups of isometries on $\mathbf{R}_+^n$ which are isomorphic as groups to the fundamental groups of M_1 and M_2 . These fundamental groups are isomorphic to each other by assumption, and so Γ_1 and Γ_2 are isomorphic to each other.

Isomorphic as abstract groups, that is. That does not necessarily mean that they sit inside of the group of all hyperbolic isometries on $\mathbf{R}_+^n$ in the same way. Mostow's theorem implies that they do. The isometry between M_1 and M_2 comes from an isometry on $\mathbf{R}_+^n$, and this implies that Γ_1 and Γ_2 are actually conjugate to each other within the group of isometries on $\mathbf{R}_+^n$. This can fail when $n = 2$ or for discrete groups of isometries which do not correspond to compact manifolds, although the compactness assumption can be weakened.

This rigidity phenomenon is actually part of a vast area in mathematics, involving other kinds of groups and geometries. See [GP] for a survey.

There is an amusing aspect of Mostow's original proof. It is not so difficult to make the groups Γ_1 and Γ_2 correspond under a mapping from $\mathbf{R}_+^n$ onto itself, the problem is that this mapping need not be an isometry. This mapping can be taken to be nice and smooth on $\mathbf{R}_+^n$, but it is more interesting to look at its behavior at the boundary hyperplane. At the boundary the mapping need not be smooth a priori, but it does satisfy a geometric condition called "quasisymmetry", which means that it distorts relative distances in a bounded way. It turns out that this condition implies the existence of differentials at almost all points on the boundary, differentials that are non-degenerate. This fact can be seen as a relative of the differentiability almost

everywhere of Lipschitz mappings. (It is important here that our initial dimension be larger than 2, so that the dimension of the boundary hyperplane be at least 2.) The existence of a well-behaved differential at a point on the boundary provides a bit of asymptotic flatness which can then be “blown up” to the whole space using the invariance of the problem under our groups Γ_1 and Γ_2 . More precisely one can show that our mapping on $\mathbf{R}_+^n$ which we know cooperates with the groups Γ_1 and Γ_2 but is not necessarily an isometry can be deformed into a mapping which does both. This uses also the compactness of M_1 and M_2 to ensure that the groups Γ_1 and Γ_2 are large enough that they both “see” all of $\mathbf{R}_+^n$, and in particular they can detect the point of asymptotic flatness on the boundary.

This outline is vague but it brings up an interesting point. In the middle of the proof one uses a theorem about differentiability almost everywhere to get the existence of a certain kind of limit even though the final statement does not really call for such a limit. One might say that it is a more infinite process than the statement seems to need. This is particularly true when one thinks about starting with finite (smooth) hyperbolic manifolds, which are more like “finite” complexity. The mapping that we create at the boundary is more like “infinite” complexity, although it has a huge amount of symmetry coming from the way that it interacts with the groups Γ_1 and Γ_2 . In the step where we obtain differentiability at some point we forget about the extra symmetry for a moment and simply treat the mapping as a general one.

Thus *this* method, which was the original one, does seem to leave the “category” of complexity of the original statement in a rather clear and interesting way.

See [GP] for other approaches to rigidity.

5.2 Groups of polynomial growth

Let G be a finitely generated group with generating set S , say. Let $V(n)$ denote the number of elements of G which can be expressed as a word of length $\leq n$ in the generating set S . We say that G has *polynomial growth* if there are constants $C, d > 0$ such that

$$(5.4) \quad V(n) \leq C n^d \quad \text{for all } n \leq 1.$$

This property does not depend on the particular choice of (finite) generating set S .

Gromov [Gr2] proved that a finitely generated group of polynomial growth contains a nilpotent group of finite index. The proof was more transcendental than one might expect the statement to require. Roughly speaking one builds certain auxiliary “continuous” spaces on which the group acts by isometries. These auxiliary continuous spaces are obtained naively from the group as limits of discrete spaces. The isometry groups of these auxiliary continuous spaces are shown to be Lie groups (with only finitely many components) through general results of Montgomery and Zippin. This method permits one to obtain a lot of homomorphisms into Lie groups, which is an important part of the argument.

In this case the statement of the theorem is already pretty concrete. The assumption of polynomial growth entails an infinite process, but it is not so bad. The method of the proof relies on processes which are much more transcendental.

5.3 A product expansion for $\sin z$

Consider the formula

$$(5.5) \quad \sin \pi z = \pi z \prod_{n=1}^{\infty} \left(1 - \frac{z^2}{n^2}\right).$$

The right-hand side makes sense because of the summability of $\sum \frac{1}{n^2}$.

This is a well-known identity from complex analysis, as on p195 of [Ah]. The proof is fairly easy to understand. We think of both sides of the equation as being functions of the complex variable z on the whole complex plane. It is a standard fact that the convergence of the product is sufficiently tame to give a *holomorphic* function on the plane. The product was chosen so that both sides of the equation vanish at exactly the same points, and so that the product is the “simplest” function with these properties. This permits us to take the quotient of the two functions to get a holomorphic function on the whole complex plane which never vanishes. The point then is to show that this quotient is actually constant. If it is constant then it must be equal to 1, because the quotient takes the value 1 at the origin.

For the proof that the quotient is constant we refer to [Ah]. Instead of the proof given on p195 I prefer to make an argument based on Theorem 8 on p207 of [Ah]. The idea is that one has some control on the behavior of our functions at infinity, in such a way that we can conclude that the logarithm

of the quotient is linear. The quotient is an even function by inspection, which permits us to conclude that the quotient is actually constant.

Thus at bottom we derive our identity from the standard miracle of complex analysis, which is that entire holomorphic functions of moderate growth are very special. These are variations on the theme of Liouville's theorem, to the effect that a bounded entire holomorphic function must be constant. The proof given on p195 of [Ah] uses a similar argument.

On the other hand we can see (5.5) as a sequence of identities about infinite series. We know the Taylor expansion of $\sin \pi z$,

$$(5.6) \quad \sin \pi z = \sum_{n=0}^{\infty} (-1)^n \frac{(\pi z)^{2n+1}}{(2n+1)!}.$$

One can also compute the Taylor coefficients of the right side of (5.5), simply by multiplying out the product and collecting terms to get the Taylor series expansion about the origin.

Of course (5.5) is equivalent to the equality of the two Taylor expansions. The individual equalities between Taylor coefficients are not quite "finite" statements in mathematics, since they involve π and infinite sums, but they are fairly concrete. They have natural finite approximations, and all of the relevant series converge absolutely.

Thus again we have a situation where the mathematical content of the statement is arguably much less "infinite" or abstract than the proof through complex analysis. This case is much easier to understand than the story of groups of polynomial growth.

5.4 Transversality

The following is a version of "Sard's theorem".

Theorem 5.7 *Suppose that U is an open subset of some $\mathbf{R}^m$ and that $f : U \rightarrow \mathbf{R}^n$ is smooth. Set*

$$(5.8) \quad \mathcal{C} = \{x \in U : \text{rank } df_x < n\}.$$

Then the image of $\mathcal{C}$ under f has Lebesgue measure zero in $\mathbf{R}^n$.

Here df_x denotes the *differential* of f at x , in the sense of advanced calculus.

See [Fe, Mi2] for proofs and more information. Note that the theorem is more subtle when $n < m$ than when $m = n$.

A particularly useful consequence of this theorem is that $\mathbf{R}^n \setminus f(\mathcal{C})$ is everywhere dense, and this was proved earlier by Arthur Brown. See [Mi2, MiS] for further information about the history of this result.

In topology one sometimes uses these results for the sake of *transversality*. Here is an example.

Theorem 5.9 *Suppose that M and N are smooth manifolds of dimension m and n , respectively, with $n \leq m$, and suppose that $f : M \rightarrow N$ is a smooth mapping. Then $f^{-1}(p)$ is a smooth embedded submanifold of M for “almost all” $p \in N$.*

As usual, “almost all”, meaning all but a set of (Lebesgue) measure zero, makes sense for any smooth manifold, without extra structure. In order to speak about the actual *measure* of a set – and not simply whether or not the measure is zero – requires additional structure, such as a volume form, or a Riemannian metric.

This theorem follows easily from the preceding one and the implicit function theorem. For a discussion of the latter in the context of smooth manifolds see [W].

This is a kind of “transversality” theorem. It is a basic tool in topology for demonstrating the existence of manifolds or submanifolds with prescribed properties while not *constructing* them exactly, and certainly not through an explicit finite recipe. For instance, it plays a role in the assertion that if M is a compact manifold of dimension m and $0 \leq k \leq m$ is an integer which satisfies $m < 2k - 1$, then the *rational* cohomology of M in dimension k is generated by the Poincaré duals of *smooth* submanifolds of M of dimension $m - k$ whose normal bundles are *trivial*. See the discussion on p232-233 in [MiS]. (It is important to exclude torsion in this discussion.)

The celebrated work of Thom [T] provides an analysis of when a given homology class is represented by a smooth submanifold, or when a given manifold can be realized as the boundary of another. These results rely heavily on transversality arguments, but of a slightly different nature from the ones above. Given a smooth mapping $f : M \rightarrow N$ and a smooth submanifold Y of N , one is interested to know if f is *transverse* to Y , or can be made so after small perturbation, and such *transversality* would imply that $f^{-1}(Y)$ is a smooth submanifold of M of the correct dimension. See [MiS] for more information.

In this way one might obtain smooth manifolds or classes of smooth manifolds with certain properties, but in terms of *complexity* this method has the unfortunate feature that it provides little information of a quantitative nature. In the context of Theorem 5.9, for instance, one might be able to get bounds on the *mass* of the submanifolds obtained through the *co-area theorem*, as in [Fe], which provides bounds on the “average” mass of $f^{-1}(p)$ for $p \in N$, but in general one might be hard pressed to do better. Even for that one would need bounds on the size of the differential of f , and they might not be so easy to obtain.

For a broader discussion of geometric bounds in the context of algebraic topology see [Gr1]. Indeed it was Gromov who first pointed out to me that there are basic questions of complexity to be addressed concerning the cobordism theory of Thom, and also the role of complexity and Lipschitz bounds for homotopy classes of mappings between manifolds (especially in connection with classical results of Serre).

Note that these issues can be discussed in the “finite” contexts of polyhedra instead of manifolds. For a combinatorial version of Theorem 5.9, for instance, see p235 of [MiS]. In this case transversality can be seen as finite and combinatorial.

5.5 Homeomorphisms

Suppose that P and Q are finite polyhedra. One can think of them as being finite unions of simplices in some Euclidean space, or in purely finite terms, characterized by the combinatorics of the intersections of the sub-simplices. There is a natural notion of *combinatorial equivalence*, in which we ask that there be a correspondence between the simplices inside P and Q which respects the way that they meet in faces, or that this be true after we *subdivide* P and Q . The effect of allowing subdivisions is to treat a single line segment and the union of two segments which meet exactly in an endpoint as being the same, for instance.

One can think of combinatorial equivalence geometrically in terms of the existence of a homeomorphism between P and Q which is *piecewise-linear*. For this we use the linear structure from the ambient Euclidean space, and the fact that simplices are truly flat.

There was a famous question (the “Hauptvermutung”) in topology about whether two finite polyhedra which are homeomorphic are actually combina-

torially equivalent. This is a very nice question, because general homeomorphisms allow infinite processes, while combinatorial equivalence is finite. As a practical matter a positive answer would be useful because it would allow one to show that combinatorial invariants of polyhedra lead to topological invariants. Classically one was interested in the topological invariance of homology groups defined combinatorially, but this was resolved in another way.

A counterexample to this question was found by Milnor [Mi1]. A more striking example appeared later from work of Edwards [E], who found exotic triangulations of spheres. See also [C1, C2, Dm1]. For this we take Q to be a standard polyhedral representation of the 5-sphere, e.g., as the boundary of a 6-dimensional simplex. The statement then is that there is another 5-dimensional finite polyhedron P which is homeomorphic to $\mathbf{S}^5$ but not piecewise-linearly equivalent to Q . This polyhedron P is obtained in a specific way, as the “double suspension of a homology 3-sphere”, which forces any homeomorphism between P and Q to undergo enormous stretching. More precisely, there is a particular polygonal curve Γ in P whose complement in P is not simply connected, and the image of this curve in Q under any homeomorphism must have “Hausdorff dimension at least 3”, as pointed out in [SS]. The latter condition implies in particular that if $h : P \rightarrow Q$ is a homeomorphism, then for sufficiently small $\epsilon > 0$ we cannot cover $h(\Gamma)$ by fewer than about ϵ^{-3} balls of radius ϵ . By contrast, for Γ itself we can use about ϵ^{-1} such balls, since it is polygonal (a finite union of straight line segments).

This implies that a homeomorphism from P onto Q cannot even satisfy a Hölder condition like (2.3) in Section 2.1 unless the exponent α is at least $1/3$. It is not known whether there are homeomorphisms in this case which are Hölder continuous of any positive exponent α .

So what does this mean in terms of finite versus infinite processes in mathematics? What would people have thought earlier in the century when they were debating about foundations and mathematical logic if they knew about *this*?

A slightly different type of situation occurs in dimension 4. There are compact smooth manifolds of dimension 4 which are homeomorphic to each other but not diffeomorphic. Because these manifolds are smooth we can realize them as finite polyhedra in a way that is compatible with their smooth structures. These polyhedra are not exotic like Edwards’ example; locally they are equivalent to standard (flat) 4-space. If the smooth manifolds were

diffeomorphic to each other then the corresponding polyhedra would be combinatorially equivalent. In this context they are not combinatorially equivalent, and the homeomorphisms between them must distort distances in a strong way. The necessity of unbounded distortion is not as easy to see in this situation as for Edwards' example. It is shown in [DoS] that the homeomorphisms cannot be "bilipschitz" or even "quasiconformal". The latter means that even *relative* distances have to be strongly distorted. (See [DoS] for details.) It is not known whether there exist homeomorphisms with bounded distortion of relative distances in the context of Edwards' example and its extensions.

The existence of homeomorphisms in these examples of 4-dimensional manifolds is provided by the work of Freedman [FQ], whose constructions entail very complicated infinite processes.

This type of phenomenon does not occur in dimensions less than 4. For dimensions ≤ 3 it is true that the existence of a homeomorphism between two smooth manifolds implies the existence of a diffeomorphism. In dimensions larger than 4 it can again happen that compact smooth manifolds are homeomorphic but not diffeomorphic or even piecewise-linearly equivalent. (See [KS, Si].) In dimensions ≥ 5 however there will always then exist a homeomorphism which is "bilipschitz", and therefore does not distort distances by more than a bounded amount. This was proved by Sullivan [Su].

Thus for *manifolds* of dimensions ≥ 5 there are still infinite processes involved but they are apparently much milder than the ones in dimension 4. Still their precise nature remains unclear though.

These examples from topology are very striking. Their geometric concreteness – finite polyhedra! – lead us to hope for similar concreteness for the associated mappings, but this simply is not true.

What are we to make of this then? Are these homeomorphisms so bizarre that we should reject them as being too far from concreteness? What would people have thought earlier in the century?

Unfortunately we do not understand so well how these homeomorphisms look. A better picture of the infinite processes involved, of natural finite approximations to them, or algorithmic processes for calculating them, would make them more concrete.

In the case of manifolds of high dimension there are fairly concrete examples [Si]. For the strange polyhedral spheres of Edwards and Cannon and the strange mappings and spaces in dimension 4 of Freedman the picture is much less clear. It can be easy to understand the spaces concretely, in finite

terms perhaps, but the mappings between them are another matter.

There is a lot of room for better theories of complexity in geometry and topology. A basic issue is to find natural phenomena of “finiteness” for the complexity of spaces and mappings between them. A first point for this is that there are only countably many compact topological manifolds up to homeomorphism type [CK]. This suggests that there is a possibility for a more “concrete” understanding.

There has been a lot of work done on finding geometric conditions on manifolds which imply that there are only finitely many underlying topological types. See [Ch, F2, GPW, Pe1, Pe2], for instance.

In [F1] there is a result to the effect that a certain kind of almost-homeomorphism is practically as good as a homeomorphism, at least in high dimensions. This type of result is very attractive if one wants to find “finite” formulations of otherwise infinite notions in topology. (One should be careful about the details here, about the quantifiers. Depending on how one formulates a notion of almost-homeomorphisms, how much one is willing to assume, it may be more or less easy to show that they can be deformed into homeomorphisms. If one makes very strong assumptions this can be achieved through a simple compactness argument. The result in [F1] is much better than that.)

The aforementioned finiteness and countability results involve somewhat similar considerations (e.g, notions of spaces being almost homeomorphic).

Alternatively one can try to find better ways to manage infinite degrees of complexity, in particular to measure them in intelligent ways. This is more the perspective of [Se5, Se9]. The idea of infinite complexity is quite basic to analysis, even if it is not described explicitly as such. This idea fits particularly well with the discussion of Section 4.4. See [DS3] for related issues of quantitative geometry and infinite complexity (some of which are also explained in the more expository [Se5, Se9]).

The idea of complexity in geometry and topology is much broader than the particular areas mentioned here. See [Gr1], for instance.

Chapter 6

Finite constructions and their limits

The idea of continuous processes is very powerful in mathematics. As in calculus, for instance. On the other hand the reliance of mathematics on continuous processes may lead one to overlook many interesting phenomena. If one thinks in terms of continuous processes one is likely to see continuous processes.

One can see the limitations of the continuous realm in all sorts of combinatorial problems. In this chapter we try to go closer to the edge between them.

This is a murky area. We are not ready to formulate a clear general principle or anything like that. We shall often find it necessary to abuse the small amount of general language that we have. In one moment we may wish to consider the real numbers as a “continuous” object, certainly so in contrast to the integers or the rationals, while in another moment we may wish to view them as “finite”, only one dimension, the “continuous” versions to be found in infinite-dimensional spaces.

Still one can attempt some general language. Often in mathematics one is interested in objects which are somehow finite but large, a finite number of particles, a finite number of parameters. One is interested in behavior as the number goes to infinity. Sometimes there are natural limiting objects to consider, sometimes not. Roughly speaking we shall try to aim towards situations where one might wish that there were a continuous limit but there is not a natural one, or where continuous models do not capture all that one wants to analyze.

It is healthy to keep in mind that for purposes of computation or physics one is often interested in numbers which are large but “far” from infinite. Numbers of modest logarithm, for instance. The restriction to relatively “concrete” large numbers is sometimes modelled well by the idea of asymptotic behavior at infinity, and sometimes not. Sometimes one cares about what happens at the level of 10^{10} but not at the level of 10^{20} , and this can effect the relevant mathematical structure. Let us begin with an example to illustrate this point.

6.1 The Hilbert transform

Let f be a function on the real line, and let us define a linear operator H on such functions by

$$(6.1) \quad Hf(x) = \text{p.v.} \int_{-\infty}^{\infty} \frac{f(y)}{x-y} dy.$$

The existence of this integral is problematic because of the singularity, but imagine writing it as

$$(6.2) \quad \int_{|x-y|>1} \frac{f(y)}{x-y} dy + \lim_{\epsilon \rightarrow 0} \int_{\epsilon < |x-y| \leq 1} \frac{f(y) - f(x)}{x-y} dy.$$

We can subtract off the $f(x)$ in the last term because

$$(6.3) \quad \int_{\epsilon < |x-y| \leq 1} \frac{1}{x-y} dy = 0,$$

as one can easily check, using the fact that $\frac{1}{x-y}$ is antisymmetric about x . This computation implies that the principal value which defines $Hf(x)$ exists whenever f has compact support and is differentiable at x , for instance. One can do much better than this, and there are deeper results to the effect that $Hf(x)$ exists almost everywhere when $f \in L^p(\mathbf{R})$, $1 \leq p < \infty$. See [Ga, Ka, St2].

This operator is called the *Hilbert transform*. It is very basic in analysis and is connected to complex analysis, partial differential equations, and operator theory. It turns out that it actually defines an isometry on $L^2(\mathbf{R})$, modulo a constant multiplicative factor. One can see this using the *Fourier transform*, for instance. A deeper fact is that H defines a bounded linear operator on $L^p(\mathbf{R})$ for each $1 < p < \infty$. See [Ga, Ka, St2].

Thus H does not distort the size of a function too badly. This is a little surprising if one simply stares at the integral naively. For instance, what if one tries to put in absolute value signs? What about

$$(6.4) \quad \int_{-\infty}^{\infty} \frac{|f(y)|}{|x-y|} dy?$$

This is a disaster. It will be equal to $+\infty$ at almost all points x at which $f(x)$ is nonzero (at all of them if f is continuous). Because we put in the absolute values we cannot make a trick like (6.2). The analogue of (6.3) does not work now.

Thus the “cancellation” in the kernel of the Hilbert transform plays a crucial role. Without the cancellation the operator is hopeless, the integral simply diverges.

But is the bad operator really so bad? To understand this it is helpful to think about what happens when one makes discrete approximations. The fact that one has bounds for H – e.g., that it defines a bounded operator on L^p – is essentially the same as saying that discrete approximations to H satisfy uniform bounds, independently of the level of the discretization. For the bad operator, with the absolute values inside, the discretizations will become unbounded in the limit.

Let us be more precise. Imagine replacing the real line with the set X_N of numbers between $-N$ and N which are integer multiples of $\frac{1}{N}$, where N is a large integer. It is easy to define versions of H and its “bad” counterpart with the absolute values inside on the discrete model X_N . A function is now a function on X_N , and one replaces integrals and integral norms with sums. The analogue of H on X_N has L^p bounds which do not depend on N when $1 < p < \infty$. One can show that the analogous operator with the absolute values inside has operator norm on L^p like $O(\log N)$.

For computationally realistic values of N the “bad” operator is essentially bounded. This was first pointed out to me by R. Coifman, who got it from V. Rokhlin. For me this was quite startling, something to resist at first, but indeed the “bad” operator is not really so bad computationally. It can be computed quickly by a computer using wavelet-based methods, as in [BCR1].

In summary, although boundedness properties of singular integral operators correspond exactly to uniform bounds for their discrete approximations, some operators that are bad in the limit are pretty good for realistic levels of discretization.

6.2 Topology

It is very natural to ask about the *topological* equivalence of two spaces. To try to track the way that pieces are connected together if not their precise size and shape.

In some cases we can try to think about this in finite terms. For instance we can look at *finite polyhedra*, and we can look at *combinatorial* or (which is the same) *piecewise-linear* equivalence. These are finite notions which are basically topological.

We also have the usual notion of topological equivalence, defined in terms of the existence of a homeomorphism. Topological equivalence is implied by combinatorial equivalence, but the converse can fail dramatically, as we discussed in Section 5.5.

Thus in topological equivalence we have a concept which is a kind of limiting version of combinatorial equivalence but which does not reduce to the latter in the cases where it could, i.e., finite polyhedra. One can try to *make* finite versions of it for finite polyhedra, through mappings which might behave like a homeomorphism at all scales above some particular small scale. This gets to be complicated, especially if we are not specifying a rate of continuity for our mappings in advance. In the infinite world we can use the notion of continuity to accommodate all rates of continuity at once, as in Chapter 2, while in the finite world this does not work so easily.

If we are simply interested in *mappings*, rather than *equivalences*, then the story is simpler. We might view continuous mappings between topological spaces as an “infinite” version of continuous piecewise-linear mappings between finite polyhedra. The “completion” to the infinite notion behaves fairly well, in the sense that arbitrary continuous mappings between finite polyhedra can be approximated by piecewise linear mappings. This is well-known and not difficult to prove, and its analogue for homeomorphisms is far from true, as we have seen.

Consider now the notion of *homotopy equivalence*. Suppose that we are given two finite polyhedra P and Q , and a pair of continuous maps $f, g : P \rightarrow Q$. We can talk about the mappings being homotopic, so that there is a continuous mapping $H : P \times [0, 1] \rightarrow Q$ which interpolates between them, $H(x, 0) = f(x)$, $H(x, 1) = g(x)$. If we have continuous mappings $f : P \rightarrow Q$ and $\phi : Q \rightarrow P$ such that $\phi \circ f : P \rightarrow P$ and $f \circ \phi : Q \rightarrow Q$ are both *homotopic* to the identity mappings on P and Q , then we get a *homotopy equivalence* between P and Q . This defines another topological notion of equivalence

between finite polyhedra, which makes sense as well for topological spaces in general. While this notion lives a priori in the infinite world, since we are allowing continuous mappings, we could also define a “finite” version, in which we restrict ourselves to mappings which are piecewise linear. This would include the “interpolating” mappings implicit in the homotopies (like H above).

These two versions of homotopy equivalence actually define the same relation between finite polyhedra. This is well known and not hard to prove, using the fact that continuous mappings between finite polyhedra can be approximated (uniformly) by piecewise linear mappings, as mentioned above. Thus we have a simpler correspondence between the “finite” and “infinite” worlds for the notion of homotopy equivalence than we have for topological equivalence.

There are variations on this theme, in which we look at mappings which are more than continuous but less than piecewise-linear, but the matter is about the same. One could use only continuous mappings which are Lipschitz or Hölder continuous, for instance. In that case the notion of homotopy equivalence would not apply to topological spaces in general but to metric spaces, for instance.

This brings up an interesting point. Suppose that we begin with the idea of homotopy equivalence between finite polyhedra in which all the mappings are piecewise linear, and imagine that we ask ourselves naively for natural “infinite” versions. We might find ourselves lead to homotopy equivalence between arbitrary topological spaces, or to homotopy equivalence between metric spaces in which all the mappings are Lipschitz or Hölder continuous, or to something else besides. Thus we can have different “completions” of the original finite concepts. Different but rather close, enjoying natural compatibilities.

If we were to start with combinatorial equivalence and look for natural “continuous” versions, we might arrive at the notion of homeomorphic equivalence between topological spaces, or something stronger for metric spaces, involving Lipschitz or Hölder continuity, for instance. Again we get different “completions” to the infinite world. This time they are much more different, without such simple compatibilities as before.

Thus finite notions might or might not have well-behaved continuous versions, and these versions might or might not be “unique”.

(Some additional technical remarks. Let us say that two metric spaces are *bilipschitz equivalent* if there is a bijection between them which does

not expand or contract distances by more than a bounded factor. This is the same as asking that both the bijection and its inverse be Lipschitz mappings. Combinatorial equivalence between finite polyhedra implies bilipschitz equivalence. We already know that topological equivalence is strictly weaker than combinatorial equivalence for finite polyhedra, and bilipschitz equivalence lies strictly between the two. Indeed, it is possible for two finite polyhedra to be homeomorphic but not bilipschitz equivalent, as observed in [SS], using the examples from [E, C1, C2]. There are also the examples in dimension 4 now, as discussed in Section 5.5. In the other direction there are finite polyhedra which are bilipschitz equivalent but combinatorially distinct. One can get this from [Si], and this is clarified further by the general theory in [Su]. Examples of the latter phenomenon could also be obtained from [Mil], although for this it seems better to rely on the later technology of H -cobordisms and s -cobordisms (as pointed out to me by S. Cappell). The analogous question for 4-dimensional compact smooth manifolds remains open, i.e., whether bilipschitz equivalence implies smooth or piecewise-linear equivalence (which are the same in this dimension).)

6.3 Finite sets and geometry

Let us try think about geometry in a primitive way. We are given something like a metric space, or even just a set inside of some $\mathbf{R}^n$. Maybe a curve or a surface. Or maybe back to a finite world, finite sets, or graphs.

“Geometry” is a tricky word. One can make many different kinds, different interpretations. The idea of a metric space provides one interpretation. Distance and nothing more.

Suppose that we are working inside some fixed $\mathbf{R}^n$. We can approximate any compact set with a finite set. To make this precise we can use the *Hausdorff metric*. Given $A, B \subseteq \mathbf{R}^n$, set

$$(6.5) \quad D(A, B) = \max\left(\sup_{a \in A} \text{dist}(a, B), \sup_{b \in B} \text{dist}(b, A)\right).$$

Thus $D(A, B) \leq \epsilon$ means that every element of A is within distance ϵ from B and vice-versa. This defines a complete metric on the set of nonempty compact subsets of $\mathbf{R}^n$. If we restrict ourselves to sets A, B which are contained inside some fixed compact set, then the resulting space is compact.

Similar considerations apply inside any metric space.

Inside this metric space of compact subsets the collection of finite subsets is dense. This is easy to show. It provides a way to make precise the idea that compact sets are simply “limiting versions” of finite sets.

We have considered compact sets as subsets of some larger space, like a Euclidean space, but only for convenience. Gromov has given a version of the Hausdorff metric for compact metric spaces as abstract spaces, independently of an embedding. See [Gr1]. Again metric spaces made from finite sets are dense within the totality of all compact metric spaces.

But is this really the way that we want to take limits of finite sets?

Let us work with subsets of $\mathbf{R}^n$. Given a finite subset F of $\mathbf{R}^n$, let μ_F denote the probability measure which is the sum of the point masses associated to elements of F divided by their total number. Thus $\mu_F(E)$ is simply the number of elements of F which lie in E , divided by the total number of elements in F . We may permit the elements of F to have repetitions.

We can take limits of these measures to get other measures on $\mathbf{R}^n$. We shall use the following standard notion of convergence. If $\{\mu_j\}_j$ is a sequence of Borel measures on $\mathbf{R}^n$, each finite on bounded subsets of $\mathbf{R}^n$, and if μ is another such measure, then we say that $\{\mu_j\}_j$ converges to μ if

$$(6.6) \quad \lim_{j \rightarrow \infty} \int_{\mathbf{R}^n} \phi d\mu_j = \int_{\mathbf{R}^n} \phi d\mu$$

for all continuous functions ϕ on $\mathbf{R}^n$ with compact support. A well known result in functional analysis implies that any sequence $\{\mu_j\}_j$ of Borel probability measures – nonnegative measures with total mass 1 – has a subsequence which converges in this sense. The limiting measure may not be a probability measure, though. It will be nonnegative and have total mass at most 1, but the total mass could be strictly less than 1. Some mass might leak out to infinity. There are simple criteria to prevent this, such as the requirement that the measures have support all contained in a common compact set.

This notion of convergence of probability measures leads us to a different way to pass to the infinite realm with the notion of finite sets. It is not hard to show that any probability measure can arise as the limit of measures associated to finite sets as above, and in fact the same is true of subprobability measures (nonnegative Borel measures with total mass at most 1) if we allow portions of the finite sets to leak off to infinity.

Suppose now that we are given a sequence of finite sets F_j in $\mathbf{R}^n$, and assume for simplicity that they are contained in a fixed compact set. We can form the corresponding measures μ_{F_j} , which we shall denote as μ_j for

convenience. Assume that the F_j 's converge to a compact set K in $\mathbf{R}^n$ in the Hausdorff metric, and that the measures μ_j converge to a probability measure μ . We can always achieve the existence of these limits by passing to subsequences. It is not hard to show that the support of μ is contained in K .

However, the support of μ may be a proper subset of K . The measure μ encodes the locations of the points weighted according to their multiplicity, and if the sets become too sparse in some region, that part might disappear in the limit of the measures, even if it persists in the limit of the sets.

For instance, suppose that a point p lies in each F_j . Then p will certainly lie in K as well. Suppose also that p is the only element of the ball $B(p, 1)$ which lies in each F_j . If we allow points to have multiplicities, then let us ask that p always have multiplicity 1 as well. If the number of elements of F_j tends to infinity as $j \rightarrow \infty$ – the situation of primary interest for us – then p will not lie in the support of the limiting measure μ , and in fact we shall have $\mu(B(p, 1)) = 0$. (For this it matters that $B(p, 1)$ is the *open* ball with center p and radius 1.)

In some ways the idea of the limiting measure contains *more* information than the idea of the limiting set, in some ways *less*. In any case they can be quite different, unless one imposes additional conditions.

Suppose now that we have sets in $\mathbf{R}^n$ which are not finite but which are finite graphs, finite unions of line segments, say. We can take limits of them in the sense of Hausdorff limits, as above, but maybe when we do that we lose some interesting geometric information.

Imagine for instance that we have a sequence of polygonal curves Γ_j which all connect some point p to some point q . We can choose the Γ_j 's so that they approximate the line segment from p to q in the Hausdorff metric – e.g., every element of Γ_j lies within a distance of $1/j$ of that line segment – but so that the Γ_j 's are also pretty jagged, constantly going up and down. In such a way that each Γ_j has twice the length of the line segment which connects p and q , for instance.

Thus something is clearly lost in the limit, some finer information about the way that the Γ_j 's behave. In the limit it is easy to go straight from p to q , but not in any of the approximations. In fact, we could choose the Γ_j 's so that they converge to the line segment in the Hausdorff distance but have lengths tending to infinity.

Here is another example. Imagine a sequence of graphs in $\mathbf{R}^2$ which approximate a square in the Hausdorff distance. That is already a bit funny,

a sequence of one-dimensional sets converging to a two-dimensional one, but it is easy to arrange. In fact one can do it in very different ways. In one case one might approximate the square by a bunch of parallel line segments. They do not have to be connected to each other. In the limit “connections” emerge which are not at all present in the approximations. In the same way that a sequence of finite sets can approximate a curve, or a square.

On the other hand one might choose a sequence of approximations which looks like a fine mesh. A union of two collections of parallel line segments, that is, vertical and horizontal. In this way all of the “connections” within the square are approximated by ones in the graphs. Approximated in the Hausdorff metric; in terms of length the approximations may not work. Suppose that we are given two points in the square which are connected by a line segment that is neither horizontal or vertical. In order to go between small neighborhoods of the two points in the approximating graphs one would have to traverse a polygonal path that switches between horizontal and vertical at least once. The lengths of the approximating paths would differ from the optimum by a definite proportion.

There is much more geometry in the approximations than we see in the limit if we simply think in terms of Hausdorff limits or limiting measures. There are more elaborate theories of “currents” and “varifolds” which take into account additional information of this type. See [Mor, Sml]. In these theories a limit of something like graphs will still be viewed as 1-dimensional even if the limit spreads out across a square or other higher-dimensional set. These theories take into account a notion of “tangent directions” in addition to the distribution of the underlying points. Both the distributions of tangent directions and points are tracked through measures, and they can become fuzzy in the limit, in the same way that a sequence of finite sets can fill out arbitrary sets in the limit.

Still these theories capture only some aspects of the geometry. Of course we cannot always expect to be able to extend a given geometric idea to a limiting regime.

6.4 Finitely generated groups

Suppose that we have a group G with a finite number of generators $g_1, \dots, g_n$. We can form *words* out of the g_i ’s, finite strings of g_i ’s and their inverses, and all elements of the group arise in this manner. There may be many

words however which represent the same group element, and this can lead to difficult combinatorial problems. This is true even if the group is finitely presented, so that the set of relations is generated by a finite set of words. The word problem may be algorithmically unsolvable, for instance. (See [Man].)

This is a situation with interesting combinatorics encoded in finite constructions, the construction of words. What are natural limiting versions of these finite constructions? One could simply take infinite words, but that may not be so interesting after all. One might hope for a natural kind of “product integral”, at least in some cases, but this is not clear. For hyperbolic groups one has the boundary at infinity as defined in [Gr3], and more generally one can look at asymptotics of the group (in the word metric). See [Gr4]. There are also natural non-hyperbolic cases with good “boundaries”. See [FM].

In the preceding examples we often had a continuous version (or versions!) that we “wanted” to believe in, that are well-established in mathematics. In this case the matter is less clear.

6.5 Comments about symmetry and infinity

Let us accept the idea that there are vast regions of unexplored territory in the world of finite and combinatorial structures, and that transcendental/continuous mathematics is of limited scope in comparison.

It is often easier to prove theorems in the infinite world. It is often easier to use the language more efficiently there to make descriptions. As in calculus, for instance; in addition to matching a continuous idealization, it is simply easier to compute in calculus than in discrete approximations. Easier to make mathematical models of physical processes, and to find clean formulations for both questions and answers.

One can also typically find more symmetries in mathematical objects in the transcendental/continuous world. Symmetries are interesting in their own right and can facilitate the proving of theorems. We have discussed this before, in Chapter 3.

Without these kinds of symmetries it may be physically impossible to find proofs, they would be just too long to find. We may also have let ourselves be trapped by the convenience of these symmetries and continuous models.

Chapter 7

Measuring infinite complexity, I

How can we make measurements of different but infinite levels of complexity?

Ideas of infinite complexity have a prominent role in classical analysis, even if they are not described explicitly as such. Again we shall look at some specific examples.

7.1 Functions and polynomials

Let us think about functions, on the real line for simplicity. Real-valued functions.

In learning calculus the first class of functions that one might encounter are *polynomials*. These are described by finitely many parameters and can be evaluated at any given point through finite arithmetic operations. We might say that polynomials have *finite complexity*. Continuous or even smooth functions in general have *infinite complexity*. How can we be more precise about this? How can we make measurements?

In calculus we learn about *Taylor polynomials* and *power series expansions*. These ideas provide ways to try to approximate general functions by polynomials. The sine, cosine, and exponential functions have power series expansions which converge everywhere and rapidly. In some cases, like the tangent function, or algebraic functions, there are power series expansions which work locally but not globally, due to the presence of singularities.

The same phenomenon occurs for rational functions, such as $\frac{1}{1-x}$. This blows up at $x = 1$ but for $|x| < 1$ it is represented by the power series

$$(7.1) \quad 1 + x + x^2 + x^3 + \dots$$

One might argue that there is little point in using power series in this case, though, since the function can already be “computed” through arithmetic operations directly.

Another example in a similar vein is the function $\frac{1}{i-x}$, where now we allow functions with complex values and i denotes the square root of -1 . This does not have a singularity on the real line, but it does have one in the complex plane, at $x = i$. This means that the power series expansion at $x = 0$ cannot converge for $|x| > 1$, because otherwise the function would not have a pole at $x = i$. One can check this directly anyway, but the point is a general one. We might try to approximate a function by polynomials, through a power series expansion, and this can lead to trouble even when there is not a singularity on the range being considered.

On the other hand we have the famous theorem of Weierstrass, to the effect that if $f(x)$ is any continuous function on a compact interval $[a, b]$, then f can be uniformly approximated by polynomials. That is, for each $\epsilon > 0$ there is a polynomial $P(x)$ such that

$$(7.2) \quad \sup_{x \in [a, b]} |f(x) - P(x)| < \epsilon.$$

Of course the polynomial P depends on f and on ϵ . See [Ru1] for a proof.

If this works for any continuous function, why do we care so much about power series? Doesn't this already say that *all* continuous functions are practically the same as polynomials?

For the second question, the answer is yes in a sense and no in a sense. Continuous functions involve infinite processes in general, and Weierstrass' theorem provides a way to say that they can be approximated by finite processes. Neither the *nature* nor the *rate* of approximation provided by Weierstrass' theorem are very strong however.

To measure the rate of approximation we can proceed as follows. Given a positive integer d , let $\mathcal{P}_d$ denote the collection of polynomials of degree at most d . This is a vector space of dimension $d + 1$. We can ask how close our given function comes to this special subspace of the functions on $[a, b]$. We make measurements using the supremum norm, defined by

$$(7.3) \quad \|g\| = \sup_{x \in [a, b]} |g(x)|$$

when g is a function on $[a, b]$. For a given function $f(x)$ on $[a, b]$, consider the quantity

$$(7.4) \quad \text{dist}(f, \mathcal{P}_d) = \inf_{P \in \mathcal{P}_d} \|f - P\|$$

defined for each d . This measures the distance from f to the polynomials of degree at most d in the supremum norm. Weierstrass' theorem says exactly that

$$(7.5) \quad \lim_{d \rightarrow \infty} \text{dist}(f, \mathcal{P}_d) = 0$$

whenever f is a continuous function on the compact interval $[a, b]$. The converse is also true, because the uniform limit of continuous functions is necessarily continuous. (See [Ru1].)

How fast do the numbers $\text{dist}(f, \mathcal{P}_d)$ go to zero?

The answer depends on the choice of function f . The numbers $\text{dist}(f, \mathcal{P}_d)$ may go to zero as slowly as they like, depending on the “degree” of the continuity of f . This can be as “bad” as it likes. To make this precise, recall that a continuous function f on a compact interval $[a, b]$ is always *uniformly continuous*. This means that for each $\epsilon > 0$ there is a $\delta > 0$ so that if $x, y \in [a, b]$ and $|x - y| < \delta$, then

$$(7.6) \quad |f(x) - f(y)| < \epsilon.$$

For a general continuous function there is nothing that we can say about the relationship between ϵ and δ .

Suppose that f is everywhere differentiable on (a, b) . Then for each $x, y \in [a, b]$ with $x \neq y$ there is a point ξ in the open interval between x and y such that

$$(7.7) \quad f(x) - f(y) = f'(\xi) \cdot (x - y).$$

This is the *mean value theorem*. From this we conclude that

$$(7.8) \quad |f(x) - f(y)| \leq \left\{ \sup_{z \in (a, b)} |f'(z)| \right\} \cdot |x - y|$$

for all $x, y \in [a, b]$. The supremum on the right hand side may not be finite, but it will be if f' is continuous on (a, b) and extends to a continuous function on $[a, b]$.

In this case we can take δ to be proportional to ϵ , i.e., we can take

$$(7.9) \quad \delta = \epsilon \cdot \left(\sup_{z \in (a, b)} |f'(z)| \right)^{-1}.$$

In general we might not be able to choose δ to be a linear function of ϵ . For instance, if $[a, b] = [0, 1]$ and $f(x) = \sqrt{x}$, then we cannot take δ to simply be a multiple of ϵ when ϵ is very small. We have to take δ to be at least $\sqrt{\epsilon}$,

as one can check by taking $x = 0$. For general continuous functions there is no universal law for the relationship between ϵ and δ . Given any proposed relationship one can find continuous functions for which it does not work.

One can bound the rate at which $\text{dist}(f, \mathcal{P}_d)$ goes to 0 in terms of the “modulus of continuity of f ”, which means (in essence) the way that δ depends on ϵ . If f is differentiable everywhere on (a, b) and its derivative is uniformly bounded, then the asymptotic rate will be faster than for $\sqrt{x}$, and in the manner that one might expect (linear decay in the first case, and not in the second).

The relationship between the rate of decay of $\text{dist}(f, \mathcal{P}_d)$ and the modulus of continuity of f is good but not perfect. The modulus of continuity can never be better than linear, unless the function is constant; for if δ were to go to 0 faster than any small multiple of ϵ , then the derivative of the function would have to vanish everywhere. The rate of approximation of a function by polynomials can be much faster than the reciprocal of the degree however. In effect it can take into account *higher* derivatives as well. For polynomials we simply have that $\text{dist}(f, \mathcal{P}_d) = 0$ for sufficiently large d . If f can be represented by a power series that converges everywhere on the real line, then we have very fast exponential decay, i.e.,

$$(7.10) \quad \text{dist}(f, \mathcal{P}_d) = O(\eta^d) \quad \text{as } d \rightarrow \infty$$

for any $\eta > 0$, no matter how small. If we take a function like $f(x) = \frac{1}{x+i}$ (with its singularity off the real line), then on any given interval $[a, b]$ we shall have exponential decay, but the rate will depend on the interval. One can verify this directly for this particular function. (As a general recipe one can employ power series representations centered on disks in the complex plane, with the center taken to lie off of the real line, at strategically-chosen locations.)

In general the rate of decay can be related to the size of the derivatives of f , when they exist. The more derivatives which exist the better, the better for decay of $\text{dist}(f, \mathcal{P}_d)$ that is. One can also look for stronger forms of convergence, e.g., uniform convergence of the functions and of their derivatives, or one could look for approximations in other kinds of norms, integral norms for instance. These issues are thoroughly studied in analysis. This is part of *Approximation Theory*, about which many books have been written.

Another issue is the problem of being able to find approximations effectively. For the Weierstrass theorem and its cousins there are fairly simple

recipes, but there are other situations where explicit solutions are not so easy to come by. There are nonconstructive methods in mathematics for proving the existence of certain kinds of approximations, and this is an interesting phenomenon for the broader themes of this book. For Weierstrass' theorem, for instance, we can rephrase the matter as saying that the polynomials are *dense* in the (Banach) space $C([a, b])$ of continuous functions on the interval $[a, b]$ equipped with the supremum norm $\|\cdot\|$ defined above. This provides a kind of simplification in the language which permits other techniques to be applied, even if it is less convenient for producing actual approximations or measuring rates of decay.

One line of reasoning would proceed as follows. Suppose that the polynomials were not dense in $C([a, b])$. Then there would be a *bounded linear functional* on $C([a, b])$ which vanishes on the polynomials. That is, there would be a linear mapping Λ from $C([a, b])$ into the scalars (real or complex numbers, as one prefers) which is *bounded*, which means that there is a constant $M > 0$ such that

$$(7.11) \quad |\Lambda(f)| \leq M \cdot \|f\|$$

for all $f \in C([a, b])$, and which satisfies

$$(7.12) \quad \Lambda(P) = 0 \quad \text{for all polynomials } P.$$

The existence of such a Λ is a standard consequence of the *Hahn-Banach theorem*. (See [Ru2].) One then tries to show that this linear functional cannot exist.

Note that the Hahn-Banach theorem relies on transfinite induction, a very large and abstract kind of infinite process.

The next step in this kind of reasoning is to say that Λ is actually represented by integration, i.e., there is a finite measure μ (but not necessarily positive) on $[a, b]$ such that

$$(7.13) \quad \Lambda(f) = \int_a^b f \, d\mu \quad \text{for all functions } f \in C([a, b]).$$

This uses a form of the *Riesz representation theorem*. (See [Ru2].)

In the context of Weierstrass' theorem this machinery is a bit ridiculous, since there are much more direct ways to solve the problem. The ideas are illustrated better by the *Müntz-Szasz theorem*. Suppose that we are working on the unit interval $[0, 1]$, and we are given a set E of positive integers. Let

$\mathcal{P}(E)$ denote the set of polynomials which are linear combinations of constant functions and monomials x^j , $j \in E$. Under what conditions on E is $\mathcal{P}(E)$ dense in $C([0, 1])$ dense? The answer is that this happens if and only if

$$(7.14) \quad \sum_{j \in E} \frac{1}{j} = \infty.$$

This is the *Müntz-Szász theorem*. See [Ru2] for a proof, which follows the lines above and then uses complex analysis to analyze the relevant measures.

Nonconstructive methods are less satisfactory for making measurements of the level of infinite complexity, but there are circumstances in which they can be used to analyze the rate of approximation in a useful way.

7.2 Polynomials, schmolynomials

Do we really want to measure the complexity of functions in terms of approximation by polynomials? Polynomials provide a natural notion of “finite complexity”, but not the only one. For this discussion we restrict ourselves to functions defined on the real line, say, or intervals within the real line.

Instead of using polynomials we could take rational functions (with poles off the domain) as our basic “finite” objects, as we mentioned earlier. We can try to measure the complexity of functions in terms of their approximations by rational functions.

We want to begin with some building blocks. We start with the rational function $\frac{1}{1+x^2}$, and then we add two parameters as follows. Given real numbers a and b , consider the function (in x) defined by

$$(7.15) \quad \frac{b^2}{b^2 + (x - a)^2}.$$

This function looks like a kind of bump. It is easy to see that this function assumes its maximum at $x = a$ and steadily decreases as x moves away from a in either direction. The value of this function at $x = a$ is 1. As one moves away from a the function starts to become small. The rate at which it does this is measured by b . Indeed, notice that our bump function is roughly constant at the scale of b ; given any interval of length b , the function does not change very much. On intervals of length larger than b it does change more significantly, the values steadily going down to zero with each step away from a .

The rate of decay is not real fast, but still pretty fast. One can make it faster by using higher powers than 2. One might hope for an exponential rate of decay, but of course one cannot achieve that with rational functions. One could use exponentials instead, like Gaussians.

At any rate our bump function (7.15) provides a very simple method for approximating continuous functions by rational functions. Fix a $b > 0$, very small, and think of approximating a given function $f(x)$ on the unit interval $[0, 1]$ (say) in the following manner. Assume for simplicity that b is the reciprocal of a large integer, $b = N^{-1}$. We can approximate f by a function of the form

$$(7.16) \quad \sum_{j=1}^N \phi_j \cdot \frac{\frac{j^2}{N}}{\frac{j^2}{N} + (x - \frac{j}{N})^2}$$

for some choice of numbers ϕ_j . This is reasonably intelligent precisely because our bump function (7.15) looks like a bump, it is roughly concentrated in the interval $[a - b, a + b]$, and it takes roughly the value 1 there. In our case we are taking the a 's to be multiples of $b = N^{-1}$, and this means that our bump functions

$$(7.17) \quad \frac{\frac{j^2}{N}}{\frac{j^2}{N} + (x - \frac{j}{N})^2}$$

should be roughly independent of each other. That is, they should not interact too much, because they decay as one moves away from their centers.

How should we choose the coefficients ϕ_j ? An obvious guess might be to take $\phi_j = f(\frac{j}{N})$. This does not work so well, even for constant functions. There are some multiplicities that we have to take into account, but that is the only real problem. To see this define a “multiplicity function” $M(x)$ by

$$(7.18) \quad M(x) = \sum_{j=1}^N \frac{\frac{j^2}{N}}{\frac{j^2}{N} + (x - \frac{j}{N})^2}.$$

This is what we would get in the sum by taking all the coefficients ϕ_j to be 1. For our given function f let us choose the coefficients to be given by

$$(7.19) \quad \phi_j = f(\frac{j}{N}) \cdot M(\frac{j}{N})^{-1}.$$

Denote by f_N the function that results from (7.17) with these coefficients. Then we have that f_N converges to f uniformly on $[0, 1]$ as $N \rightarrow \infty$. Let us see why this is so.

Fix a point x , and consider $f_N(x)$, which is given explicitly by

$$(7.20) \quad f_N(x) = \sum_{j=1}^N f\left(\frac{j}{N}\right) \cdot M\left(\frac{j}{N}\right)^{-1} \cdot \frac{\frac{j}{N}^2}{\frac{j}{N}^2 + (x - \frac{j}{N})^2}.$$

This is a weighted average of the values $f(\frac{j}{N})$ of f . It is an *average* because the weights are positive and their sum is 1, the latter a consequence of the presence of the multiplicity function. The point now is that if N is very large then these weights are concentrated at the j 's for which $x - \frac{j}{N}$ is small compared to N^{-1} . When $x - \frac{j}{N}$ becomes large compared to N^{-1} , the weights become small, and the contribution of those terms is small. In short $f_N(x)$ is an average of values of f with the average concentrated at values of f taken at points $\frac{j}{N}$ which are very close to x when N is large enough. One can then use the (uniform) continuity of f to say that this average is (uniformly) close to $f(x)$.

One needs to make some computations to make this argument precise, but this is not very difficult. One can also see clear relationships between the modulus of continuity of the given function and the rate of approximation of the f_N 's. These can be formulated as statements about the relationship between the modulus of continuity and the rate of approximation as measured by the *degree* of the rational functions (in this case the f_N 's). The main point is that the parameter b (in this case $1/N$) corresponds to the resolution of the approximation in a way that can easily be made precise.

This is one way to make approximations by rational functions, but it is by no means the only way, or the most efficient way. There are better methods for taking into account more smoothness or more subtle oscillations, but we shall not pursue this here. One of the main points is to use rational functions and parameters like a and b to treat *locations* and *scales* simultaneously and individually.

Still this gives a flavor, and a way to look at measuring complexity of functions. Note that some functions admit much better approximations by rational functions than by polynomials. Rational functions can deal with singularities more efficiently, and our bump functions (7.15) are localized much better than polynomials ever would be.

If one likes localization, one might as well ask about approximation by step functions instead. Recall that a *step function* is simply a linear combination of characteristic functions of intervals. One might also describe them as being piecewise-constant. It is very easy to approximate a continuous

function on a compact interval uniformly by step functions, simply using the uniform continuity of the function. Step functions are also described by only a finite number of parameters, and so provide another notion of “finite complexity” from which to work.

Of course step functions are also discontinuous, which causes trouble in some contexts. One might prefer to approximate functions by others which are at least as regular as the original. Thus one might consider instead approximations by *piecewise linear functions*. For these it is reasonable to demand continuity when approximating continuous functions. For piecewise-linear functions one gives up differentiability at the break points. To get higher degrees of smoothness one can look at functions which are piecewise polynomial, like *polynomial splines*.

Another deficiency of approximation by step functions is that the rate of approximation will never be so great, never more than linear basically. By using piecewise linear functions or splines of higher order one can get faster rates of approximation of functions which have some smoothness.

The idea that we want to approximate functions by functions which are as good as the original points to a deficiency in the earlier method for approximating functions by rational functions. While it enjoyed a certain naive simplicity, it suffers from the fact that to get a good approximation we use bump functions which are highly concentrated, with the resolution parameter b chosen very small, so that our basic building blocks are relatively singular. They have large derivatives, for instance.

There are more efficient methods of approximation for avoiding this problem. The area of harmonic analysis often called “Littlewood-Paley theory” deals with this issue in a good way. Basically one should make approximations that are adapted to scale and location, and not just individual locations. Thus if a function is almost constant in some region, it is not very efficient to approximate it by highly concentrated bumps there. Now-a-days these matters are typically discussed in the context of “wavelets” (*ondelettes* in French).

One can make other methods of approximation. A particularly important one is to use *Fourier series*. Suppose that we have a function on the real line, continuous say, which for simplicity is periodic with period 1. One can try to realize it as a sum of simpler functions of the form $\sin 2\pi nx$, $\cos 2\pi nx$, where n is a positive integer. One should also include constant functions.

There are many results about being able to do this. Let us call a function a “trigonometric polynomial” if it is a finite linear combination of sines,

cosines, and constants as above. A variant of the Weierstrass approximation theorem states that continuous periodic functions can always be approximated uniformly by trigonometric polynomials. If one works instead in L^2 one gets an orthogonal basis from the trigonometric functions, so that an L^2 function can always be expressed as an infinite series of trigonometric functions with the coefficients computed by simple integrals. The series will converge in the L^2 norm, but it may not converge uniformly even if the original function were continuous. These are well-studied issues, with many results known. See [Ru2, Z], for instance.

In a certain sense the analysis of functions by trigonometric functions is very similar to analysis through polynomial approximation. One can view periodic functions as being functions on the unit circle in the plane $\mathbf{R}^2$, and trigonometric polynomials on the line correspond exactly to the usual polynomials on $\mathbf{R}^2$ restricted to the unit circle.

In other ways these various kinds of approximation and expansions are very different. One of the reasons that Fourier series are very useful is that partial differential equations, such as the equation which models the vibration of a string, become much simpler when one expands functions in terms of Fourier series. An individual sine or cosine function can represent a “standing wave”, and the frequency n is very important, corresponding to pitch for sound. These things do not work as nicely for ordinary polynomials, or for rational functions.

On the other hand Fourier series are not very good for treating local behavior in an efficient manner. If one has a function that is very calm most of the time but then erratic at a particular location, then it is hard to see that in the Fourier series, because the effect of the erratic behavior spreads out. By using rational functions, or splines, or wavelets, one can treat local phenomena differently.

One might also have a large region in which the function is very calm, another where it is not. Wavelets are pretty good at figuring out which is which, and treating them automatically. That is, they are good for accommodating calm or erratic behavior that depends on the function in an automatic way.

There is a more elaborate way to try to “analyze” the behavior of functions, based not on a (linearly independent) basis of basic functions, but on a larger collection from which one can make a library of bases. Different bases to see different kinds of behavior, be it waveform, localized oscillations, something more spread out. One then has algorithms for choosing the best basis from the library (where “best” has to be explained). See

[Co, CMQW, CW1, CW2].

In short, there are many different methods for “analyzing” functions, which lead to different ways to measure complexity.

7.3 Linear operators

Let us consider a very concrete linear operator which arises in mathematics. Let $f(x) = f(x_1, x_2)$ be a function on $\mathbf{R}^2$, and consider the operator defined by

$$(7.21) \quad Rf(x) = \text{p.v.} \int_{\mathbf{R}^2} f(y) \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} dy.$$

This is a *singular integral operator*, because the integral does not exist in the usual sense. One has to define it as a *principle value* (as in Section 6.1). This means that in fact we set

$$(7.22) \quad Rf(x) = \lim_{\epsilon \rightarrow 0} \int_{|x-y|>\epsilon} f(y) \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} dy.$$

If f is smooth and compactly supported (or even much less) one can show that this limit always exists, by rewriting it as

$$(7.23) \quad \int_{|x-y|>1} f(y) \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} dy + \lim_{\epsilon \rightarrow 0} \int_{\epsilon < |x-y| \leq 1} (f(y) - f(x)) \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} dy.$$

To make this rewriting we use the fact that

$$(7.24) \quad \int_{\epsilon < |x-y| \leq 1} \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} dy = 0,$$

which can be checked directly. (It is easiest to see this using symmetry considerations, noticing how the integrand changes sign when one reflects across an axis.) The point now is that if f is smooth and compactly supported, then

$$(7.25) \quad \left| (f(y) - f(x)) \frac{(x_1 - y_1)(x_2 - y_2)}{|x - y|^4} \right| \leq \left\{ \sup_{z \in \mathbf{R}^2} |\nabla f(z)| \right\} \cdot |x - y|^{-1}.$$

This follows from the fact that $|f(y) - f(x)|$ is no greater than $|x - y|$ times the supremum of $|\nabla f|$ on the line segment which joins x to y . Once we have this estimate (7.25) it is easy to show that the limit in (7.23) exists. (Indeed now the “singular” integral converges even if we put absolute values inside the integral, which was not true before.)

Thus the singular integral exists for nice functions f . This operator arises mathematically in the following way. Recall that the Laplacian Δ is the second-order differential operator defined by

$$(7.26) \quad \Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$$

(since we are in two dimensions). Suppose that we have a smooth function f with compact support, and we want to be able to derive information about the various second derivatives of f from the knowledge of Δf alone. A basic formula is

$$(7.27) \quad \frac{\partial^2}{\partial x_1 \partial x_2} f = R(\Delta f)$$

(except for a multiplicative constant). Thus R is really the quotient of $\frac{\partial^2}{\partial x_1 \partial x_2}$ and Δ . This formula holds for practically any function, modulo suitable interpretation.

This kind of identity can be obtained as in [St2], as well as many others like it. For the moment the main point is that this is a linear operator of basic importance in mathematics, and there are many others like it. What can we say about it in terms of complexity, and the infinite processes which it involves? Is it “simpler” than an arbitrary operator, can it be described more “efficiently” than in (7.21) above?

In this case the answer is yes. Our operator enjoys extra symmetries. It commutes with translations, for instance. This implies that it can be “diagonalized” by the Fourier transform. More precisely, if $f(x)$ is a function on $\mathbf{R}^2$, then its *Fourier transform* is the function $\hat{f}(\xi)$ on $\mathbf{R}^2$ defined by

$$(7.28) \quad \hat{f}(\xi) = \int_{\mathbf{R}} f(x) e^{-2\pi i \langle x, \xi \rangle} dx.$$

We are using here the standard inner product $\langle \cdot, \cdot \rangle$ on $\mathbf{R}^2$. This is like the definition given in Section 4.1 for functions on the line, and the inversion of the Fourier transform works in the same manner as indicated there. For our

operator R we have that

$$(7.29) \quad R(\widehat{f})(\xi) = \frac{\xi_1 \xi_2}{|\xi|^2} \widehat{f}(\xi),$$

except for a multiplicative constant. In other words, the complex exponentials are eigenfunctions for R , and they are a “complete set”. (We are ignoring some technical details here, but they are standard. See [SW] for more information about Fourier transforms and this fact that they diagonalize operators which commute with translations.)

The fact that we have a diagonalization is a way to say that there is less “complexity” in this matrix than we might have thought. A general $N \times N$ matrix has N^2 entries, a diagonal matrix has N entries. One could make suitable discretizations in the present story to get closer to matrices if one wanted to.

In fact all sorts of operators are “diagonalizable”, in the same way that real symmetric matrices can be diagonalized, by the spectral theory. (See [Ru3], for instance.) This sounds pretty good, but for considerations of complexity one should also take into account the transformation (“change of basis”) needed to effect the diagonalization. In this respect the Fourier transform behaves fairly well. It can be computed efficiently, through the fast Fourier transform.

Thus one might say that the complexity of our operator R is much less than it might appear to be at first, or than it would be for an arbitrary operator. There are still some problems with this, however. The Fourier transform works very well for some kinds of information but not others. It is not easy to see localizations in the Fourier transform. By contrast it turns out that R does behave fairly well in terms of localization. That is, the effect of R on a function in a certain region is not too sensitive to what happens far away. The Fourier transform is not like that, it is more global.

The precise localization properties of R are a bit tricky. Some of them are described in [St2]. They are reflected in part by the boundedness of R on many natural Banach spaces of functions. This is not at all clear from the diagonalization through the Fourier transform, because the action of the Fourier transform on most function spaces is not so easy to understand.

For example, R defines a bounded linear operator on $L^p(\mathbf{R}^2)$ when $1 < p < \infty$. This means that for each such p there is a constant $C(p)$ such that

$$(7.30) \quad \int_{\mathbf{R}^2} |Rf(x)|^p dx \leq C(p) \int_{\mathbf{R}^2} |f(x)|^p dx$$

for all functions f on $\mathbf{R}^2$. This kind of result is very useful in analysis, especially partial differential equations; it provides a way to control the size of the $\frac{\partial^2}{\partial x_1 \partial x_2}$ derivative of a function in terms of the size of the Laplacian. It does not work for $p = 1, \infty$, and in particular it is better to measure “size” here in terms of certain types of averages instead of the supremum. There are many other natural function spaces on which R is also bounded.

For $p = 2$ this boundedness can be proved using the Fourier transform. The main points are that the Fourier transform preserves the L^2 norm, and on the Fourier transform side R looks like the operator of multiplication by a bounded function. This does not work for other L^p spaces because the Fourier transform does not have such simple behavior on them. The fact that R does behave well on them is a reflection of its sensitivity to localization and the underlying geometry of $\mathbf{R}^2$.

There are other methods from Harmonic Analysis for analyzing operators which avoid some of the difficulties suffered by the Fourier transform. Now-a-days people often speak in terms of “wavelets”, which provide special bases for function spaces which have moderate sensitivity to both location and oscillation while at the same time having natural scale-invariance. For instance, one can characterize L^p spaces in terms of wavelets in a reasonably simple way when $1 < p < \infty$, purely in terms of the *sizes* of coefficients. This is not true for the Fourier transform when $p \neq 2$. This is called the *unconditional basis property*, and it is a good substitute for orthogonality which is meaningful in non-Hilbert spaces.

It turns out also that when we represent R as a matrix in terms of a wavelet basis it is almost diagonal in a precise sense which is useful in practice.

In fact wavelets cooperate well with a variety of linear operators and function spaces. The Fourier transform is better for getting very sharp information in very special situations, while wavelets are typically less precise but more flexible. Ideas related to wavelets are often applicable in cases where there is no translation-invariance or even a group structure necessarily. We have emphasized the operator R for the sake of illustration, but the methods are actually very general.

See [Db2, Db3, Me1, Me2] for more information about wavelets and the analysis of functions and operators. Note that they are powerful tools simply for studying the complexity of functions, independently of operators.

Many of these ideas for analyzing the complexity of functions and operators existed in Harmonic Analysis long before the more recent advent of wavelets. One of the main points about wavelets is that one can get actual

orthogonal bases with these good properties; before one had various methods for breaking functions and operators into simple pieces and putting them back together again, and these methods enjoyed much the same properties in terms of localization in space and frequency, but the orthogonality properties were not as exact. Orthogonality is often very convenient even in situations where its exact form is not crucial.

Wavelets also work well for numerical computation. In part this comes from their orthogonality, which eliminates some of the redundancies needed as well as providing exact formulas that do not require additional analysis. Another important point is that the wavelets themselves can be computed efficiently, and the representations of functions in terms of wavelets can be computed efficiently. This uses the particular structure of the wavelets defined by Daubechies in [Db1]. It is not that there is a simple formula for these wavelets, but rather an iterative construction which lends itself to numerical computation. For applications to the efficient computation of operators like R , see [BCR1, Co].

7.4 General comments about operators

Let us look a bit more about general issues of “infinite complexity” related to linear operators which arise naturally in mathematics.

Notice that an operator like R can be discretized in “meaning” as well as in the formula for its definition. One could work on a discrete 2-dimensional lattice, define discrete versions of the Laplacian and the mixed second derivatives, and look for a “quotient” as before (as in (7.27)). One would still get something which looks approximately like R .

Thus in a sense the concept of R is not unavoidably infinite. The “infinite” realm is more convenient for certain types of theoretical considerations, though.

There are other aspects of operator theory that do require the infinite world in a stronger way. For this discussion I wish to stick to fairly “concrete” operators, of the sort which arise in connection with partial differential equations, as opposed to more abstract operator theory.

In the “pseudodifferential calculus” one has a collection of linear operators which includes standard differential operators but which also allows more complicated constructions. One has a way to try to take inverses, square roots, and other operations of interest.

Actually, one does not perform these operations in an *exact* way, but instead one has a symbolic calculus for converting questions about operators into simpler questions about functions, at the expense of allowing certain kinds of errors. The composition of two operators corresponds to the multiplication of a pair of functions (the “principal symbols” of the operators), but one can recover the operators from the principal symbols only to first approximation, modulo an error which is of “lower order”. One can make more refined calculations to make the errors even more modest, but the improvements are in terms of *smoothness* rather than *size*.

In other words, by allowing certain types of errors one can make symbolic calculations which detect much of the mathematically significant information in a convenient manner. One has a good theory here of the “infinite” part of the complexity.

For theoretical purposes this can be quite powerful. One might be interested in tracking singularities, for instance, or one might not mind being able to invert an operator only up to a finite rank error. In some cases such ambiguities are unavoidable.

For the matter of smooth or finite rank errors we see a very clear difference between infinite mathematics and finite approximations. For infinite mathematics finite rank errors are not so bad, but in a discrete approximation or numerical computation of actual solutions the matter is far different. See [BCR2] for a treatment of pseudodifferential calculus in a computational setting.

Similarly the distinction between *smoothness* and *singularity* reflects an important difference between finite and infinite situations. In the mathematical study of partial differential equations which describe the evolution of some object, like a fluid, one is very interested in knowing about the possibility and nature of “blow-up”. In finite realms the distinction is much less sharp.

For the question of invertibility, one of the basic points behind this discussion is the following. Suppose that we have an operator T acting on some Banach space, and that we know that it is a perturbation of a simpler operator T_0 that we understand fairly well already. Let us write E for the error, so that $T = T_0 + E$. If we know that T_0 is invertible, then under what conditions on the error term E can we invert T ?

This is true if E is sufficiently small (in operator norm), by a well-known computation. In the context of differential equations, or the pseudodifferential calculus, one typically knows instead that the error term is somehow

smooth instead of small. In practice this leads to some form of *compactness*, like the statement that E can be approximated by finite rank operators in the operator norm. Thus we can proceed to $T = T_1 + E_1$, where T_1 is a small perturbation of T_0 and therefore invertible, while E_1 is now of finite rank.

Given the knowledge that T is a finite rank perturbation of an invertible operator, in order to show that it is invertible it suffices to show that it is either injective or surjective. This is true automatically in finite dimensions but not in general in infinite dimensions.

In the context of partial differential equations this fact can be very useful. The problem of inverting an operator might correspond to the problem of solving a differential equation with some boundary conditions, for example. The preceding facts can permit one to reduce a question of existence to one about uniqueness, for instance, which might be approachable by direct calculation.

Thus in the end the mathematical theory sometimes clears away the brush until one comes down to a finite problem. The mathematical theory often stops then.

There is another aspect of “infinite complexity” in operator theory, in which one has an operator in hand with some compactness properties and one wants to analyze more precisely the way that it can be approximated by finite-rank pieces. This issue is analogous to the one of approximating functions by simpler pieces, and indeed there are various ways in which the two questions are closely connected. See [Smb, RS], for instance.

Chapter 8

Measuring infinite complexity, II

We want to continue looking at mathematical tools for measuring infinite complexity, but now with more emphasis on geometry and topology.

8.1 Sizes of sets

In the finite realm one might simply measure size by counting the number of points, but for infinite sets the matter is more complicated. We certainly want to say that a plane is bigger than a line even if they are the same in terms of cardinality.

Consider a subset E of some $\mathbf{R}^n$. How can we measure its size? The following is a simple way. For each $\epsilon > 0$ let $N(\epsilon)$ denote the smallest number of balls of radius ϵ that it takes to cover E . This will be finite as soon as E is bounded. One then looks at the asymptotic behavior of $N(\epsilon)$ as $\epsilon \rightarrow 0$. For instance, if

$$(8.1) \quad N(\epsilon) = O(\epsilon^{-d}) \quad \text{as } \epsilon \rightarrow 0$$

for some $d \geq 0$, then this is a way to say that E is at most d -dimensional in terms of size.

For example, a line segment satisfies (8.1) with $d = 1$, and a square satisfies (8.1) with $d = 2$. For an entire line $N(\epsilon)$ is always infinite, as it is for any unbounded set. In that case it is better to restrict oneself to bounded

subsets and make measurements there, and then look to see how the bounds depend on the subset, e.g., on its diameter.

In this kind of measurement the “dimension” parameter d need not be an integer. For instance, for the usual Cantor set the correct d is $\log 2 / \log 3$.

Of course one is not restricted to taking powers in (8.1), one can use more complicated functions too.

This method of measuring size has some disadvantages. Suppose that E is a countable set whose closure contains a ball. Then the best that one can do with $N(\epsilon)$ is say that it satisfies (8.1) with $d = n$, which is true of any bounded subset of $\mathbf{R}^n$. This is not so nice, and one might want to say that such sets are very small. For this one has to look more deeply into the structure of the set. Let us describe a couple of ways for doing this.

Given $d \geq 0$, set

$$(8.2) \quad A^d(E) = \inf \left\{ \sum_j (\text{diam } X_j)^d : \{X_j\}_j \text{ is a sequence of subsets of } \mathbf{R}^n \text{ which covers } E \right\}.$$

This is called the d -dimensional *Hausdorff content* of E . It differs from the above because one can look at coverings of variable size, i.e., the X_j 's do not have to all have the same diameter. For instance, $A^d(E) = 0$ whenever E is countable and $d > 0$. For something like a line segment or a square the Hausdorff content would coincide with the usual measure, modulo a normalizing factor, and assuming that one took the right dimension d .

A problem with the Hausdorff content is that it is always finite, at least for bounded sets. A square has finite 1-dimensional content, for instance; the finiteness of $A^d(E)$ does not really say something about E , in terms of the dimension d . By contrast, a condition like (8.1) does see the dimension in a strong way, (8.1) is not true for a square with $d = 1$.

The Hausdorff content can see the dimension in a different way. If S is a square then $A^d(S)$ is always finite, but it is 0 as soon as $d > 2$. More generally, if a set E satisfies (8.1) for some d , then one has $A^s(E) = 0$ for all $s > d$.

In the Hausdorff content the X_j 's do not have to be small. By contrast the asymptotics of $N(\epsilon)$ as $\epsilon \rightarrow 0$ must unavoidably reflect the behavior of the set at small scales. We can do this also for Hausdorff content through a more precise definition.

Let $d \geq 0$ and $r > 0$ be given. Again d corresponds to dimension, while

r will correspond to a length scale. Set

$$(8.3) \quad H_r^d(E) = \inf \left\{ \sum_j (\text{diam } X_j)^d : \{X_j\}_j \text{ covers } E \text{ and} \right. \\ \left. \text{diam } X_j < r \text{ for all } j \right\}.$$

In other words, $H_r^d(E)$ is the same as $A^d(E)$ except that all the X_j 's are required to be small. It is easy to see that $H_r^d(E)$ increases as r decreases, because one is then taking infima over smaller sets. Set

$$(8.4) \quad H^d(E) = \lim_{r \rightarrow 0} H_r^d(E).$$

This defines the d -dimensional *Hausdorff measure* of E . The limit always exists because of the monotonicity in r , but we allow the value ∞ .

This is now more sensitive to the dimension. If S is a square then $H^2(S)$ is the same as the usual 2-dimensional measure of the square, except perhaps for a normalizing constant, but $H^d(S) = \infty$ when $d < 2$, as it should be. Similar remarks hold for line segments or the standard Cantor set, with respect to the correct dimension in each case.

If E satisfies (8.1) for some d , then $H^d(E) < \infty$ for that choice of d . The converse is not true. Countable sets have 0 H^d -measure for each $d > 0$, but they can be more complicated for (8.1). One does not even have to take countable dense sets, simply a sequence covering to a point will work if the convergence is not too fast. Similarly one can look at sequences of line segments, or little squares.

Indeed, one of the main advantages of Hausdorff measure and Hausdorff content over measurements as in (8.1) is that they are *countably subadditive*; the size of a countable union is less than or equal to the sum of the measures. One of the main advantages of Hausdorff measure over Hausdorff content is that Hausdorff measure is *countably additive*, the measure of a countable disjoint union of *measurable* sets is the sum of the individual measures. The failure of countable (or even finite) additivity of Hausdorff content reflects the fact that it is not required to see small scales. When $d = 0$, for instance, H^d simply counts the number of elements in the given set, while A^d can be much smaller.

On the other hand countable additivity does allow “messy” sets with many little pieces floating around. In some situations one might want to forbid that, and measurements in terms of $N(\epsilon)$ can be more appropriate. It depends on the situation.

Notice that $H^d(E) = 0$ if and only if $A^d(E) = 0$. We have

$$(8.5) \quad A^d(E) \leq H^d(E)$$

for all sets, just from the definitions, but the reverse inequality is wrong, as we have seen. However, $H^d(E) = 0$ when $A^d(E) = 0$, because in that case the efficient coverings of E for computing $A^d(E)$ had to involve sets of small diameter.

The *Hausdorff dimension* of E is defined by

$$(8.6) \quad \inf\{d \geq 0 : H^d(E) = 0\}$$

Note that $H^d(E) = 0$ for all $d > n$ when E is contained in $\mathbf{R}^n$, and so the Hausdorff dimension is no greater than n in this case, which is good. One can also talk about the (*upper*) *Minkowski dimension* as the smallest d for which (8.1) holds, although for unbounded sets it is better to look at the smallest d such that (8.1) holds for all bounded subsets. There are other notions of dimensions and other measures related to these, as in [Fa, Fe, Mat].

Sets in Euclidean spaces can be complicated. Very complicated. We have many ways to measure how complicated they are. So far we have talked about size, but indeed also the geometry in a slightly more subtle way, through the structure of the coverings, the different ways that coverings can be organized. These coverings speak about the geometry of a set, maybe in a tricky way if we allow coverings by sets of different diameters, as in Hausdorff measure and Hausdorff content.

These ideas can make sense in “finite” contexts, in discretizations, but it is not completely easy to do something beyond simply computing the number of elements in a set. One should distinguish between scales in the discretization, e.g., by working with a version of H_r^d in which r is much larger than the smallest scales of the discretization but still very small. Otherwise, if one simply takes r to be the minimal scale of the discretization, one cannot do more than simply count points.

Even this can be useful, especially if one makes additional localizations, e.g., counting the number of points in the set inside balls of certain radius. Looking at this as a function of the radius, comparing the result with its counterpart for the whole space, etc.

Still it is much easier to distinguish between different scales in infinite mathematics than in finite worlds.

Even for questions of size and covering, it is not really clear that we have settled on good measurements of “complexity”.

8.2 Complexity of sets

How might we measure complexity of sets?

There are many answers to this question. Here is one. Suppose that we have a set in some $\mathbf{R}^n$ which is somehow “1-dimensional”. In terms of Hausdorff measure for instance. We can then ask whether it is contained in a rectifiable curve, and if so how long the curve should be.

The constraint that a set be contained in a rectifiable curve imposes some nontrivial conditions on its size, but that is not the only matter. There is also a question of how *scattered* the set is.

Let us consider a concrete example, a Cantor set in the plane constructed in the following manner. One starts with the (closed) unit square and then takes the four corner squares of one-fourth the size. For each of these squares one does the same, and so forth. At the k^{th} stage of the construction one has 4^k squares of size 4^{-k} . One takes the union of these 4^k squares at the k^{th} stage to get a closed set E_k , and then one takes the intersection of all the E_k ’s to get a Cantor set E which has dimension 1 in all the senses mentioned in the previous section. In particular, it satisfies (8.1) with $d = 1$, just like a line segment.

In terms of covering properties this set has practically all the same properties as a line segment. However one can show that it is not contained in any curve of finite length, or even in a countable union of them. (See [Mat].)

If a set is contained in a rectifiable curve, then one can use the length of the shortest such curve as a measurement of its complexity. A measurement of “infinite complexity”, because it still allows infinitely many corners, loops, etc.

To be contained in a countable union of rectifiable curves imposes already nontrivial restrictions on structure. Roughly speaking this property says that the set has approximate tangent lines at almost all points. See [Fa, Fe, Mat] for details. The Cantor set described above is far from having approximate tangent lines.

In [J2, O] there is a characterization given of the compact sets K which are contained in a curve of finite length, with bounds for the length of the optimal curve. This characterization is given in terms of certain quantities which measure how close K lies to a line inside a given ball. One looks at balls at all scales and locations and combines them in a certain way to get a number which turns out to be comparable to the length of the shortest curve which contains K . See [J2, O] for a more precise statement and further

details. A key point is to take *orthogonality* into account, in this case through the Pythagorean theorem.

This story is related to the earlier one about functions. We can measure the complexity of a function through its oscillations, like the “modulus of continuity” mentioned before. Now it is more natural to measure the complexity of a function in terms of the degree to which it can be approximated by affine functions. This is the analogue of measuring the complexity of a 1-dimensional set by the extent to which it can be approximated by a line segment at different scales and locations, and it is closely related to the topics in Section 4.4.

For either functions or sets one makes measurements in arbitrary balls, say, and then combines all the measurements for all balls to get some global measurement of complexity. This is a very basic theme in harmonic analysis, and it has natural geometric counterparts as well.

In this regard functions and sets are closely related. If the structure of a set is to be controlled through something like a “parameterization”, as in the idea of putting a set inside a rectifiable curve, then the parameterization involves functions whose complexity as functions will be related to the geometric complexity of the sets. This will not necessarily work for all natural measurements of the complexity of the functions, and not all of them will have natural geometric meaning.

See [DS1, DS3, Se5, Se9] for more information about these themes, and also for analogous matters in higher dimensions. For the “qualitative” versions (allowing countable unions of surfaces) see [Fa, Fe, Mat] also for higher dimensions.

One should keep in mind that these are all measurements of *infinite* complexity. Even when a quantity like the length of a curve is finite, *infinitely* many corners or loops or crossings are permitted. This is true of the other measurements that we have in mind here as well. The point is to control the “level of infinity”.

Of course these are not the only ways to think about complexity of sets. These are measurements of complexity in comparison with *Euclidean geometry*, broadly construed. Self-similar sets can also be considered as “nice”. They can admit rapid computational description, for instance. Sometimes approximations of given sets by self-similar ones are even used for the purposes of image compression. See [Ba].

The difference between roughly Euclidean geometry and models which are self-similar but fractal is a bit like our earlier stories for functions and

operators. One can make measurements of complexity based on different methods of analysis – the Fourier transform versus wavelet bases, for instance – and these can lead to very different answers.

8.3 Topological dimension

Let us now consider a different measurement of complexity of sets, namely *topological dimension*. The basic reference is [HW]. For simplicity we shall restrict our attention to compact metric spaces, but the theory works more generally.

The definition of topological dimension in [HW] is inductive. One starts by saying that a space has topological dimension 0 if it is totally disconnected. A space has dimension $\leq n$ if each point has a system of neighborhoods whose boundary has dimension $\leq n-1$. A space has dimension n if it has dimension $\leq n$ but not $\leq n-1$.

I prefer another characterization. A compact metric space $(M, d(x, y))$ has dimension $\leq n$ if for each $\epsilon > 0$ there is a finite collection of open sets $\{U_i\}$ in M such that the U_i 's cover M , $\text{diam } U_i < \epsilon$ for all i , and no point in M lies in more than $n+1$ of these sets.

For example, an interval has dimension ≤ 1 but not dimension < 1 unless it is a point. This is not hard to check. The fact that a square has dimension ≤ 2 is slightly more amusing. One has to be a bit careful in the way that one makes the covering, but it is not difficult.

It is important to notice what this characterization of topological dimension does not ask for. It does not ask for a bound on the total number of the U_i 's. In effect the condition controls the local structure of the space but not at all how large it is.

The condition also does not impose any requirements on the shape of the U_i 's, only that they be sufficiently small.

Of course the condition is genuinely *topological*, in the sense that it does not depend on the choice of metric, only on the underlying topology. Change the metric and the space might be much “larger” in terms of size, but not in terms of the topological dimension. For instance, if the original metric is $d(x, y)$, and we change it to $d(x, y)^{\frac{1}{2}}$, then this is still a metric – the main point is that the triangle inequality continues to hold – and dimensions which measure size (like the Hausdorff dimension) increase by a factor of 2 under this change, but the topological dimension remains the same. Similar results

hold if one replaces $\frac{1}{2}$ with any number s so long as $0 < s < 1$.

There is a nice theorem of Alexandroff, to the effect that a compact metric space M has topological dimension $\leq n$ if and only if it can be approximated by finite polyhedra of dimension $\leq n$ in the following sense. Given an arbitrary $\epsilon > 0$ one asks for a finite polyhedron P and a continuous mapping $f : M \rightarrow P$ with the property that each “fiber” $f^{-1}(p)$, $p \in P$, has diameter $< \epsilon$. This does not imply that M is homeomorphic to a subset of finite polyhedron of dimension $\leq n$, but it wants to approximate that statement. For instance, suppose that M is a subset of a plane, the Cartesian product of the unit interval $[0, 1]$ and a totally disconnected infinite compact set, like a convergent sequence together with its limit. It is easy to see that this set has topological dimension ≤ 1 , but of course it is not homeomorphic to a subset of a 1-dimensional polyhedron.

Note well that the polyhedron P is permitted to depend on ϵ . One can also look at its complexity, total number of faces, for instance. The notion of topological dimension does not impose any restriction on the complexity of P , except for its dimension, but one can profitably impose such conditions. See [Se8], for instance.

Another nice theorem says that if M has topological dimension $\leq n$ then M can be homeomorphically embedded into $\mathbf{R}^{2n+1}$. Thus the notion of topological dimension provides a characterization of the compact metric spaces which can be embedded into some finite-dimensional Euclidean space.

This embedding result is quite remarkable, in the way that it involves infinite processes. The direct part of Alexandroff’s theorem, the passage from the bound on the topological dimension to the polyhedral approximation, is a simpler matter of finite combinatorics and partitions of unity. One fixes an $\epsilon > 0$, applies the hypothesis at a particular scale, obtains the conclusion at a particular scale. By contrast the embedding theorem involves distances at all scales; one has to build a mapping for which the inverse image of a point is a point, and not merely small in diameter. The usual proof relies on the Baire category theorem.

The embedding theorem provides a nice example of a result which involves infinite processes and for which it is convenient not to make the proof at all explicit!

Another way to understand topological dimension involves the notion of *stable values*. Let M be a compact metric space, and let I_n denote the closed unit cube in $\mathbf{R}^n$. We say that a point p in I_n is a *stable value* of a continuous mapping $f : M \rightarrow I_n$ if there is a $\delta > 0$ so that if $g : M \rightarrow I_n$ is any

continuous mapping which is δ -close to f , i.e.,

$$(8.7) \quad |f(x) - g(x)| < \delta \quad \text{for all } x \in M,$$

then p lies in the image of g . Of course this implies that p lies in the image of f , and we call $p \in f(M)$ an *unstable value* of f if it does not enjoy this property.

It turns out that if M has topological dimension strictly less than n then every value of every continuous mapping $f : M \rightarrow I_n$ is unstable, and that if M has topological dimension at least n then there is a continuous mapping $f : M \rightarrow I_n$ with a stable value. (One can take the stable value to be the center of the cube I_n .)

In other words, if the topological dimension is strictly less than n , then M is too “thin” to have a mapping into I_n which is very “serious”. Think about Peano curves, which can provide examples of continuous mappings from a line segment onto a square. Or the Cantor function, which maps the Cantor set continuously onto an interval. It is easy to map spaces of lower dimension onto spaces of larger dimension, but these mappings can be approximated by mappings with much smaller image. This is very easy to see for the examples just mentioned, and indeed they are typically obtained as the limit of such mappings.

The fact that spaces of dimension at least n admit mappings to cubes of dimension n with stable values is very nice. It is a strong way of saying that the space really has some n -dimensional flesh. This result requires infinite processes, and the usual proof uses the Baire category theorem.

There are other results to this effect, that having topological dimension $< n$ means that the space is approximately “invisible” for topological considerations of dimension n or greater, in terms of homotopy properties of mappings into spheres, and in terms of homology and cohomology. See [HW] for details.

One might think of the *topological dimension* as being analogous to the notion of *depth* as a measurement in complexity theory. Let us think about making constructions on a fixed polyhedron P of dimension n , constructions of a continuous mapping with some properties, say. A common line of reasoning begins by making choices at the vertices, then the edges, then the 2-dimensional faces, and so forth through dimensions until one reaches the top dimension n . It happens sometimes that the constructions on the different faces of a given dimension are more-or-less independent of each other,

that the more tricky point is the number of layers, the “depth”, which in this case means the dimension. Thus the complexity of the polyhedron – the total number of constituent simplices for instance – is sometimes not so crucial, while the dimension is. Although this is simpler for polyhedra than general spaces, similar considerations apply in general, with Alexandroff’s approximation theorem providing a convenient bridge.

This type of phenomenon comes across nicely in [Pe2]. See also [Se8].

The Brouwer whose name appears so frequently and prominently in [HW] is the same as the eminent logician.

8.4 Topological dimension, size, and structure

The topological dimension is a very good measurement of “size” of a space for looking at the behavior of continuous mappings and other topological questions. In other ways it provides practically no information; metrically the space could be very large. We can make this precise with the concept of Hausdorff measure and Hausdorff dimension. We defined these in Section 8.1 only for subsets of Euclidean spaces, but the definitions extend to metric spaces in a straightforward way.

A metric space can have topological dimension 0 but large Hausdorff dimension, or even infinite Hausdorff dimension. One can make versions of the usual Cantor set construction within spaces of larger dimension, or even infinite-dimensional spaces, to get examples. One can also proceed more abstractly. Suppose that $(M, d(x, y))$ is a metric space of Hausdorff dimension d . We can apply the *snowflake transform* to it to get the space $(M, d(x, y)^s)$, where $0 < s < 1$. This is again a metric space (the triangle inequality remains valid), and it has Hausdorff dimension d/s .

Of course the snowflake transform does not change the underlying topological structure, or the topological dimension. It is irrelevant for purely topological considerations such as the behavior of continuous mappings, the existence of stable values, etc. And yet it affects the *geometry* of the space in a dramatic way. In terms of how many balls of a given radius are needed to cover the space, for instance.

In this regard topological dimension is rather remarkable for extracting a *topological* measurement of size which does not worry too much about

geometric measurement of size.

Is there a relationship between these measurements of size? Is the geometric one always larger than the topological one, for instance?

The answer is yes. The topological dimension is always less than or equal to the Hausdorff dimension. Note well that the topological dimension is necessarily an integer, while the Hausdorff dimension need not be.

This result can be found in [HW]. In fact there is a more precise statement: if M has topological dimension at least n , then M has *positive* Hausdorff measure in that dimension as well. (These results in [HW] for Hausdorff dimension and positive Hausdorff measure are actually stated in terms of Hausdorff *content*, but this is equivalent, as in the remarks of Section 8.1.)

What about a converse? Topological dimension need not control Hausdorff dimension for a *particular* metric, but topological dimension is independent of the choice of metric. Can one always find a metric which is compatible with the topology and for which the Hausdorff dimension is equal to the topological dimension? The answer is yes, and one can even embed a given compact metric space of finite topological dimension into a finite-dimensional Euclidean space in such a way that the image has Hausdorff dimension equal to the topological dimension, where Hausdorff dimension is now computed using the ambient Euclidean metric. See [HW].

It may not be possible to find a metric on a given space so that the space has *finite* Hausdorff measure in the topological dimension. One can make easy counterexamples using the Cartesian product of an interval with an infinite set.

In summary, topological dimension and Hausdorff dimension both provide measurements of the structure of a set. They are both measurements of “infinite” levels of complexity; they tolerate infinite complexity but control it in some ways and not in other ways. They provide very different measurements of complexity.

The ideas of Section 8.2 point to another measurement of complexity still. A measurement that looks for a *geometric* resemblance to Euclidean spaces rather than a topological one, or mere similarity in terms of rough measurements of size.

Hausdorff dimension is always at least as big as topological dimension, what happens when they are equal? The topological dimension provides a kind of “floor” for the geometric size, but if the actual size is not much larger, then can we expect some nontrivial restrictions on the geometric structure? Results which permit one to obtain stronger conclusions about the geometric

complexity of the space? There are some results of this nature, and this general question is discussed in [Se5].

8.5 Lipschitz mappings

A *Lipschitz mapping* between metric spaces is one which does not expand distances by more than a constant factor.

More formally, let $(M, d(x, y))$ and $(N, \rho(u, v))$ denote two metric spaces. We say that a mapping $f : M \rightarrow N$ is *Lipschitz* if there is a constant $C > 0$ so that

$$(8.8) \quad \rho(f(x), f(y)) \leq C d(x, y) \quad \text{for all } x, y \in M.$$

We might say C -Lipschitz to make the constant more precise.

It will be convenient to use the phrase *Lipschitz function* to mean a Lipschitz mapping into the real line (with the standard metric).

The Lipschitz condition provides a measurement of infinite complexity of functions. It allows infinite amounts of activity, even for functions on the unit interval, but nonetheless imposes nontrivial restriction on this activity.

To put this measurement into perspective it is helpful to compare the Lipschitz condition with our earlier discussion of complexity of functions, and to view our earlier discussion with new eyes of suspicion. The Lipschitz property has the nice feature that it is very geometric and simple, making sense for mappings between arbitrary metric spaces. Most of the earlier discussion was not so robust, requiring much simpler structure for both domain and range. Think of approximations by polynomials, rational functions, splines, trigonometric polynomials, etc. All of these require more precise structure of the underlying space even to be formulated.

Continuous (real-valued) functions can always be approximated by Lipschitz functions, modulo some technical hypotheses. Let $(M, d(x, y))$ be a metric space, and suppose for a moment that M is compact. Let $f : M \rightarrow \mathbf{R}$ be a continuous function, and set

$$(8.9) \quad f_j(x) = \inf\{f(y) + j d(x, y) : y \in M\}.$$

This defines a real-valued function on M , and one can show that f_j is j -Lipschitz. This is not hard, using the fact that $x \mapsto d(x, y)$ is 1-Lipschitz for each $y \in M$, which itself comes from the triangle inequality. One can also prove that f_j converges to f uniformly on M as $j \rightarrow \infty$. This is well known

and not very difficult to verify, and we omit the details. The main point is to use the fact that f is uniformly continuous since it is continuous and M is compact. Indeed one can relate the rate of convergence of $f_j \rightarrow f$ to the modulus of continuity of f in a simple way.

This argument works on general metric spaces if one makes some additional assumptions. If M is not compact then the finiteness of f_j is not assured, but it can be verified under moderate assumptions on f (i.e., its growth at infinity). Once one knows that f_j is finite at any point one can show that it is finite everywhere and j -Lipschitz. Uniform convergence to f can be proved under the assumptions of uniform continuity of f and restrictions on the growth of f at infinity. (Boundedness would be plenty.)

Under modest conditions on the complexity of M one can make other procedures of approximation which are linear in f and more obvious “locality” properties. This is discussed in [Se9].

Lipschitz mappings cooperate well with our measurements of *sizes* of sets. Let $(M, d(x, y))$ and $(N, \rho(u, v))$ be two metric spaces, and suppose that $f : M \rightarrow N$ is C -Lipschitz. Let E be any subset of M . Then the size of E can be related to the size of $f(E)$ in the following ways. Given a dimension $d \geq 0$, we have that

$$(8.10) \quad H^d(f(E)) \leq C^d H^d(E) \quad \text{and} \quad A^d(f(E)) \leq C^d A^d(E).$$

This is easy to check from the definitions. Similarly, if $N(E, \epsilon)$ is defined as in the beginning of Section 8.1, as the smallest number of balls of radius ϵ needed to cover E , then we have that

$$(8.11) \quad N(f(E), C\epsilon) \leq N(E, \epsilon).$$

A particular consequence of these inequalities is that Lipschitz mappings never increase Hausdorff dimension. However they can increase topological dimension. To see this, let us first define a “symbolic” version of the Cantor set. We take K to denote the set of all infinite binary sequences, i.e., all sequences $\{x_j\}_{j=1}^\infty$ such that $x_j \in \{0, 1\}$ for all j . We equip this set with a metric by setting $D(\{x_j\}, \{y_j\}) = 0$ when $x_j = y_j$ for all j ,

$$(8.12) \quad D(\{x_j\}, \{y_j\}) = 2^{-l} \quad \text{when } x_j = y_j \text{ for all } j \leq l, x_{l+1} \neq y_{l+1}.$$

It is easy to see that this defines a metric on K in such a way that K is compact. One can also check that K has Hausdorff dimension 1.

This metric is substantially different from the one on the standard middle-thirds Cantor set which is induced by the usual Euclidean metric on the line, because they give different Hausdorff dimensions. The standard metric for the middle-thirds Cantor set corresponds approximately to a power of this metric.

Define a mapping $f : K \rightarrow [0, 1]$ by

$$(8.13) \quad f(\{x_j\}) = \sum_{j=1}^{\infty} x_j 2^{-j}.$$

One can check that this mapping is 1-Lipschitz and also surjective. However, K has topological dimension 0, while $[0, 1]$ has topological dimension 1. Thus Lipschitz mappings can indeed increase the topological dimension.

Of course Lipschitz mappings can decrease topological dimension, e.g., by mapping $[0, 1]$ to a point. One cannot map $[0, 1]$ into a set like K in an interesting way, the image will always have to be a point. In situations like this – continuous mappings between compact spaces – in order for the topological dimension of the image to be less than that of the domain it is necessary that there be a point in the range whose inverse image has topological dimension at least the difference between the topological dimensions of the domain and image. See Theorem VI 7 on p91 of [HW].

8.6 Differentiability, rectifiability, etc.

There are issues of infinite complexity for Lipschitz mappings which are much more subtle than mere size. A Lipschitz function on a Euclidean space is differentiable almost everywhere. Thus one does not simply have bounds on the sizes of the oscillations of Lipschitz functions, but also a kind of asymptotic rigidity at almost all points. See [Fe, St2, Se9]. A well-known variant of this famous result says that given any $\epsilon > 0$ one can find a C^1 function which agrees with the given Lipschitz function except on a set of measure $< \epsilon$. (See [Fe].)

There are also quantitative versions of these rigidity properties of Lipschitz functions on Euclidean spaces, as discussed in [Do, DS3, Se5, Se9]. (See also Section 9.4 below.)

To put these matters into perspective, consider the *Hölder continuous* functions on $\mathbf{R}^n$. That is, $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is *Hölder continuous of order α* if

there is a constant $C > 0$ so that

$$(8.14) \quad |f(x) - f(y)| \leq C |x - y|^\alpha$$

for all $x, y \in \mathbf{R}^n$. This is the same as the Lipschitz condition when $\alpha = 1$. When $\alpha > 1$ it is too strong, and implies that f must be constant. When $\alpha < 1$ it is weaker than the Lipschitz condition in the way that it controls the local behavior of f . One can see that there is nothing like the above rigidity properties of Lipschitz functions for Hölder continuous functions of order $\alpha < 1$. This is discussed in more detail in [Se9].

Similarly, for general metric spaces one cannot hope to have the kind of rigidity properties for Lipschitz functions that one has on Euclidean spaces. Lipschitz functions on our symbolic Cantor set K of Section 8.5, for instance, do not enjoy such rigidity properties. Nor do Lipschitz functions on Euclidean spaces equipped with “snowflake” metrics of the form $|x - y|^s$, $0 < s < 1$. (A Lipschitz function on such a snowflake space is essentially the same thing as a Hölder continuous function with respect to the ordinary Euclidean metric.) There are some metric spaces which are far from Euclidean in their geometry but for which one still has a version of differentiability almost everywhere, namely Heisenberg groups with their Carnot metrics. See [Pa].

Thus Lipschitz functions are very interesting in terms of infinite complexity. They cooperate with measurements of size in an obvious way, but they can also enjoy additional rigidity properties which are not explicitly part of their definition. These rigidity properties do not work for metric spaces in general, but depend on the geometry of the underlying space. These “hidden” rigidity properties are absent also for many other conditions that one might impose on functions, like Hölder continuity.

Just as one had differentiability (almost everywhere) and other natural “rigidity” properties of functions, there are geometric conditions like *rectifiability* and *uniform rectifiability* for sets. In the geometric setting one loses much of the linear structure of functions and methods for analyzing them (like the Fourier transform and wavelets), but one gains a more rich language, so that there are completely different types of criteria for the presence of good structure. Criteria involving the relationship between size and topology, for instance. See [Fa, Fe, Mat, DS3, Se5, Se9].

For further discussion about the analogy between functions and sets and the methods for analyzing them see [DS2, DS3], and note the differences in their conclusions!

See [DS4] for a search for other types of rigidity properties related to Lipschitz mappings and geometric measure theory.

The preceding paragraphs are very sketchy, little more than pointers to a sizable literature. In short the main point is that in these directions there are worlds of situations in which infinite complexity arises naturally and nontrivial ways to measure it and to distinguish between different levels of complexity.

8.7 Wild embeddings

Given an arc in a Euclidean space, can we always straighten it by a global homeomorphism?

By an *arc* we mean a set which is homeomorphic to the unit interval $[0, 1]$. We might say “topological arc” for emphasis. The question asks whether we can find a global homeomorphism from the ambient Euclidean space onto itself which sends the arc to a straight line segment.

In $\mathbf{R}^1$ this is trivially true because all arcs are line segments. In $\mathbf{R}^2$ this is true but nontrivial, one of the standard facts about plane topology. Starting in $\mathbf{R}^3$ it fails to be true. See [Moi], for instance.

Similarly, given a compact set in $\mathbf{R}^n$ which is homeomorphic to the usual Cantor set, one can ask whether there is a global homeomorphism on $\mathbf{R}^n$ which takes the given set to a subset of a line. Again the answer is yes in dimension 2 and no starting in dimension 3. Basic examples are provided by “Antoine’s necklaces”, which are topological Cantor sets in $\mathbf{R}^3$ with non-simply connected complement. If there were a global homeomorphism carrying such a set to a subset of a line, then the complement would have to be simply-connected (because the set is totally disconnected).

The construction of the Antoine’s necklace is easy enough to describe. For the usual Cantor set one takes a closed interval, replaces it by two disjoint closed subintervals, performs a similar operation on those, etc. For the Antoine’s necklace one starts with a solid torus and replaces it by a chain of solid tori contained inside. These new solid tori are disjoint but linked. In each of these solid tori one repeats the construction, and does this forever. In the limit one gets a set which is homeomorphic to a Cantor set, but the complement is not simply connected, as one can verify. See [Moi] for details.

Similarly there are homeomorphic embeddings of the 2-sphere into $\mathbf{R}^3$ which are “wild”, i.e., cannot be transformed into a standard 2-sphere by a

global homeomorphism. See [Moi] again, and also [BuC, Dm1].

In all of these situations one has a subset of a Euclidean space which is homeomorphic to something fairly simple but for which the embedding has to have an infinite amount of twisting. These phenomena also lend themselves naturally to questions of complexity. One can for instance look at finite versions of the constructions and measure the amount of distortion needed for global homeomorphisms which make the straightenings. The impossibility of straightening the infinite constructions implies the absence of uniform bounds for the finite versions. (Uniform bounds on the moduli of continuity of the relevant homeomorphisms and their inverses, that is.)

8.8 The Whitehead continuum

Is every contractible manifold homeomorphic to a ball?

In dimensions 1 and 2 this is true and classical. It is not true starting in dimension 3. The first counterexample was obtained by taking the complement of the *Whitehead continuum* in the 3-sphere. Roughly speaking, the *Whitehead continuum* is defined as follows. One starts with a (compact) solid torus T_1 , which we think of as embedded in the 3-sphere. That is our first approximation. For the second approximation we take another solid torus T_2 which is embedded inside T_1 . The embedding is rather special; T_2 is stretched around the middle of T_1 and has its ends clasped together on the other side. (See [Dm2, Ki] for pictures.) This clasping makes it impossible to shrink T_2 to a point without leaving T_1 and without having a copy of T_2 cross itself in the course of the shrinking. On the other hand T_2 can be shrunk to a point inside T_1 if one does allow copies of T_2 to cross themselves in the shrinking (but still not to ever leave T_1). This is because of the way that T_1 is stretched around the center of T_2 ; it does not simply go around in a circle, and in particular does not contain a homotopically nontrivial loop in T_1 .

T_2 is only the second approximation to the Whitehead continuum. For the third we embed a new solid torus inside T_2 in the same way that T_2 was embedded into T_1 . This procedure is repeated indefinitely, at each stage a new solid torus is embedded into the last one in the same way that T_2 was embedded into T_1 . To get the Whitehead continuum one takes the intersection of this decreasing chain of embedded tori. Let us call the final result W . This is a compact set, and it is easy to see that it is connected.

In fact, $\mathbf{S}^3 \setminus W$ is *contractible*. This means that it can be continuously

deformed inside itself until it becomes a single point.

$\mathbf{S}^3 \setminus W$ is also an open subset of $\mathbf{S}^3$, hence a (noncompact) manifold without boundary. It is contractible but not homeomorphic to a 3-dimensional open ball. The reason is that it is not *simply-connected at infinity*. To explain what this means it is helpful to call an open subset of $\mathbf{S}^3 \setminus W$ a *neighborhood of infinity* if it is the complement in $\mathbf{S}^3 \setminus W$ of a compact subset of $\mathbf{S}^3 \setminus W$. If $\mathbf{S}^3 \setminus W$ were simply-connected at infinity it would mean that for each neighborhood U of infinity there is a smaller neighborhood V such that every loop in V can be contracted to a point inside U . This is not true for $\mathbf{S}^3 \setminus W$. See [Dm2, Ki].

Similar issues arise in the wild examples mentioned in the previous section. Sometimes the problem is more blatant, e.g., a region is not even simply connected when it “should” be, like the complement of Antoine’s necklaces. In all cases there is an “asymptotic” problem, a problem with homotoping loops in the complement of the set but which lie near the set.

8.9 Decompositions

Imagine taking $\mathbf{R}^3$ and some compact set $K \subseteq \mathbf{R}^3$ and shrinking K to a point while leaving the other points alone. What happens to the topology?

If K is a straight line segment the resulting space is *homeomorphic* to $\mathbf{R}^3$. The operation of collapsing the line segment to a point is pretty severe geometrically, but it does give back the same topological space.

This is not true if K is a circle. If we collapse a circle to a point then the complement of that point in the resulting space is the same topologically as the complement of the circle in $\mathbf{R}^3$, and is not simply-connected. Since the complement of a point in $\mathbf{R}^3$ is always simply-connected, the new space that we made cannot be homeomorphic to $\mathbf{R}^3$.

If we take K to be the Whitehead continuum (taking care to have it inside $\mathbf{R}^3$ and not just $\mathbf{S}^3$), then the resulting space is again different from $\mathbf{R}^3$ topologically. Indeed, every point in $\mathbf{R}^3$ has arbitrarily small punctured neighborhoods which are simply connected, and this is not true when we collapse the Whitehead continuum to a point. This corresponds to the failure of simple-connectivity at infinity described in the preceding section.

Similarly if one collapses a wild arc to a point one may not get $\mathbf{R}^3$ back again.

In these cases one not only does not get $\mathbf{R}^3$ back again, one does not even get a topological manifold, because of the behavior near the strange point.

More generally one can collapse families of compact sets to points rather than just individual ones. One can simply speak of *decompositions* of $\mathbf{R}^3$ (or other spaces), in which one takes a partition (by closed sets say) and uses this to make a quotient topological space. This is discussed in more detail in [Dm2]. There are natural conditions to impose to avoid pathologies as one gets by collapsing a circle or a Whitehead continuum to a point, but there is not a simple way to tell when one will get back $\mathbf{R}^3$ or something strange.

What about natural finite approximations to making spaces by collapsing subsets to points? Can one make measurements of complexity? One way to do this is given in [Se6]. There one considers decompositions obtained through a certain kind of iteration, for which the Whitehead continuum is a simple special case. The compact sets to be collapsed are obtained as limits of compact domains obtained by repeated embeddings (roughly speaking). This kind of construction has a natural *topological* “self-similarity”, to which one associates a natural geometry which is self-similar in the usual sense. In this geometry the compact sets to be collapsed to points have diameter zero, as they should, but their approximations are shrinking at a particular rate. The final spaces can be viewed as subsets of $\mathbf{R}^4$, with the induced geometry.

In making these geometries it makes sense to stop the construction after finitely many generations. These approximations are automatically homeomorphic to $\mathbf{R}^3$, the question is what kinds of estimates one can have for the homeomorphisms, in terms of the approximately self-similar geometry. Some observations and examples are given in [Se6], but these quantitative issues have yet to be studied much.

8.10 Domains in Euclidean spaces

Let Ω be an open subset of some $\mathbf{R}^n$. Let us define the *quasihyperbolic metric* on Ω to be the Riemannian metric given at an arbitrary point $x \in \Omega$ by the formula

$$(8.15) \quad \frac{ds^2}{\text{dist}(x, \partial\Omega)^2}.$$

Thus the length of an infinitesimal bit of curve at x is the same as the ordinary Euclidean length divided by the distance to the boundary of Ω .

This defines a metric which blows up as one goes to the boundary. Not only does the infinitesimal form blow up, but the length of any path which goes toward $\partial\Omega$ will blow up. The approximate geometry is captured well by the following observation. In the geodesic distance for the quasihyperbolic metric, a ball with center $x \in \Omega$ and radius 1 is approximately like a Euclidean ball with center x whose radius is proportional to $\text{dist}(x, \partial\Omega)$ by a positive constant that stays away from 1 by a definite amount.

One might just as well define the quasihyperbolic metric using a smooth function $\delta_\Omega(x)$ instead of $\text{dist}(x, \partial\Omega)$, but where $\delta_\Omega(x)$ is chosen so that it is bounded from above and below by positive multiples of $\text{dist}(x, \partial\Omega)$ (as on Theorem 2 on p171 of [St2]).

If the complement of Ω is compact one might prefer to adjust this definition, working on the sphere instead so that the point at infinity is included, etc. Of course one can extend this construction to domains on smooth manifolds.

Using this metric one can look at the (Euclidean) boundary behavior of Ω as asymptotic geometry at infinity with respect to the quasihyperbolic geometry. If Ω is bounded by a compact smooth hypersurface, say, then the quasihyperbolic metric looks a lot like the usual models for hyperbolic geometry. Otherwise it can be much more complicated.

This geometry is often natural for analysis. Bounded harmonic functions on Ω , for instance, have very tame behavior on quasihyperbolic balls of bounded radius, while their behavior near the boundary can be much more complicated.

If the boundary of Ω is itself complicated, the quasihyperbolic metric provides a natural way to think about exploring it. Imagine that Ω is a sea, and that we are in a small boat exploring the fjord which borders it. In the examples of Section 8.7 there is interesting behavior already on bounded subsets of Ω with respect to the quasihyperbolic metric. There can be loops which cannot be filled by disks within a bounded region (in the quasihyperbolic metric), and this behavior can be repeated infinitely often as one approaches the boundary of Ω . One way to make interesting measurements is to count the number of times that such events occur, e.g., how numerous they are within regions of a certain size. This idea works well in the context of “uniformly rectifiable sets” and “almost flat hypersurfaces” (Section 9.3), for which there are typically infinitely many “complications” whose frequency can be counted with fairly good precision. See [DS1, DS3, Se1, Se2, Se3, Se4, Se9].

Alternatively one can look at all locations in Ω in a uniform way but try to deal in a more subtle way with the range and nature of the complications. See [Al, AV1, AV2, HY, V].

8.11 Spaces of bounded geometry

Let (M, g) be a complete Riemannian manifold of dimension n . Let us say that (M, g) has *bounded geometry* if for each $x \in M$ the ball in M with center x admits a diffeomorphic mapping onto an open subset of $\mathbf{R}^n$ in such a way that this diffeomorphism distorts distances by only a bounded factor and its derivatives are uniformly bounded. We do not allow these bounds to depend on the point x , but the bounds on the derivatives are permitted to depend on the level of differentiation.

In a space of bounded geometry the local behavior is controlled in a strong way, and the issue is what happens at infinity.

Examples 8.16 (1) Euclidean and hyperbolic spaces have bounded geometry.

(2) Homogeneous spaces always have bounded geometry.

(3) Domains in Euclidean spaces equipped with their quasihyperbolic metrics (as in Section 8.10) are spaces of bounded geometry, at least if one takes the trouble to replace $\text{dist}(x, \partial\Omega)$ in (8.15) by a smooth version of the distance, as discussed above.

(4) The universal covering of any compact Riemannian manifold (without boundary) has bounded geometry.

(5) A leaf of a smooth foliation in a compact manifold has bounded geometry (through the induced metric).

These examples are in a sense very different from each other, and they have lives of their own, but they can be profitably seen to some extent as special cases of a general notion. They also make for good examples of infinite complexity within ordinary mathematics. There are many natural ways to measure this asymptotic complexity, including isoperimetric inequalities. Some references include [Gr1, Gr3, Gr4, At1, At2, AtH].

Chapter 9

Some results about geometric complexity

In the preceding chapters we saw a range of situations in which one can make measurements of complexity even when the complexity is infinite. In this chapter we discuss a few particular results of this nature.

9.1 Discrete parameterizations

The following result is proved in [Sto].

Theorem 9.1 *Let r and s be positive integers, with $s \geq r$. Then there is a bijection $f : \mathbf{Z}^r \rightarrow \mathbf{Z}^s$ such that*

$$(9.2) \quad |f(x) - f(y)| \leq C |x - y|^{\frac{r}{s}}$$

for some constant $C > 0$ and all $x, y \in \mathbf{Z}^r$.

This is quite nice. We are all familiar with space-filling curves, we know that they have to be somehow “fractal”, but that is not at all the same as getting the discrete version to work out right, on the nose.

The Hölder exponent $\frac{r}{s}$ in (9.2) is absolutely the right one. It exactly matches the differences in the “dimensions” of $\mathbf{Z}^r$ and $\mathbf{Z}^s$. Notice, for instance, that the number of elements of $\mathbf{Z}^r$ which lie in a ball of radius $L > 1$ centered at the origin is bounded from above and below by constant multiples of L^r , and there is a similar statement for $\mathbf{Z}^s$. One can derive the sharpness of the Hölder exponent $\frac{r}{s}$ from this observation.

To put the theorem into perspective it is helpful to think more about its continuous analogues, mappings from $\mathbf{R}^r$ to $\mathbf{R}^s$. For this one cannot ever get bijectivity when $s > r$, by standard results in topology. Bijectivity can only occur in a discrete setting, but then what does it mean exactly?

One can think of bijectivity as a way to say “measure-preserving”, for instance. It implies the following “regularity property”, which arose first in [Dv1]. If f is as in the theorem, then there is a constant $K > 0$ so that for each ball B of radius t in $\mathbf{R}^s$ we can cover the inverse-image $f^{-1}(B)$ by $\leq K$ balls of radius $t^{\frac{s}{r}}$. This observation is analogous to one in [Dv1], and can be proved as follows. Let t and B be given as above. We may as well assume that $t \geq 1$. Let A be any subset of $f^{-1}(B)$ with the property that $|x - y| \geq t^{\frac{s}{r}}$ for all $x, y \in A$ with $x \neq y$. We want to derive a uniform bound for the possible number of elements in A . Given $x \in A$, let $B(x)$ denote the ball $B(x, \frac{1}{2} t^{\frac{s}{r}})$. The balls $B(x)$, $x \in A$, are pairwise disjoint, and so their images are pairwise disjoint as well, since f is injective. On the other hand their images $f(B(x))$, $x \in A$, are all contained in the ball obtained by dilating B by a constant factor (which depends on f but not on B). This is easy to check, since we know that $f(x)$ lies in B for each $x \in A$, and since the Hölder condition on f implies that each $f(B(x))$ has diameter bounded by a constant multiple of t . We know that each ball $B(x)$ contains about $(t^{\frac{s}{r}})^r = t^s$ elements of $\mathbf{Z}^r$ (to within a constant factor), and so the number of elements of $\bigcup_{x \in A} f(B(x))$ is at least a constant multiple of t^s times the number of elements of A . All these points must lie within a fixed dilate of B , and this implies that A contains no more than a bounded number of elements, with a bound that does not depend on B . This implies the bounded covering property that we want.

The measure-preserving and regularity properties make sense for mappings between $\mathbf{R}^r$ and $\mathbf{R}^s$ as well. In fact, the “discrete” result of Theorem 9.1 implies continuous versions through normal families arguments, as follows. Fix r and s as above, and choose f as in Theorem 9.1 and so that $f(0) = 0$. We can always arrange this normalization. Given $\epsilon > 0$, let f_ϵ be defined by

$$(9.3) \quad f_\epsilon(x) = \epsilon^{\frac{r}{s}} f\left(\frac{x}{\epsilon}\right),$$

so that f defines a bijection between $\epsilon \cdot \mathbf{Z}^r$ and $\epsilon^{\frac{r}{s}} \cdot \mathbf{Z}^s$. Each f_ϵ satisfies the analogue (9.2), with the same constant, because we were careful about the scaling. The Arzela-Ascoli argument permits one to find a sequence of ϵ 's tending to 0 such that the corresponding f_ϵ 's “converge uniformly on compact subsets” to a mapping $g : \mathbf{R}^r \rightarrow \mathbf{R}^s$. There is a small technical

problem here, since the f_ϵ 's do not all have the domain, but this is not a serious problem, because the domains become increasingly thick. This kind of issue is discussed in some detail in [DS4]. The mapping g that results might be considered as a continuous version of the mapping f from Theorem 9.1, since it can be derived from f in this way.

Our limiting mapping g also satisfies the analogue of (9.2), since the f_ϵ 's do, and with a uniform bound. One can also check that g is *regular* in the sense described above, i.e., if B is any ball in $\mathbf{R}^s$ of radius t , then $g^{-1}(B)$ can be covered by a bounded number of balls of radius $t^{\frac{s}{r}}$. This can be derived from the fact that the f_ϵ 's satisfy the same condition with uniform bounds.

The regularity condition is a kind of uniform and scale-invariant way to say that g has only bounded multiplicities. It does not imply injectivity in general. So what does the injectivity of the mapping f from Theorem 9.1 really mean? What does it imply at the continuous level?

From regularity we also get that g does not distort measures very much. That is, there is a constant M so that

$$(9.4) \quad M^{-1} |E| \leq |g(E)| \leq M |E|$$

for all, say, *compact* subsets of $\mathbf{R}^r$. Here $|E|$ and $|g(E)|$ denote the Lebesgue measures of the subsets E and $g(E)$ of $\mathbf{R}^r$ and $\mathbf{R}^s$, respectively. We do not really care about compactness of E here, it is just a convenient way to avoid measurability problems. One could as well work with outer measures and have a similar estimate for *all* subsets E of $\mathbf{R}^r$. The upper bound in (9.4) follows from the fact that g is Hölder continuous, while the lower bound can be derived from the regularity property. (For this it is convenient to use the description of Lebesgue measure as Hausdorff measure of the correct dimension. Indeed, (9.4) has a more general version for mappings between metric spaces and their effect on Hausdorff measures. This is spelled out in Lemma 12.3 of [DS4].)

These bounds on the distortion of measure fit well with the “measure-preserving” property of bijections between discrete sets, but they are less precise. They permit bounded multiplicities, like foldings, for instance. Consider instead the condition that

$$(9.5) \quad |h(E)| = |E|$$

for all compact subsets E of $\mathbf{R}^r$, where $h : \mathbf{R}^r \rightarrow \mathbf{R}^s$? This is much stronger, closer to injectivity. It implies that disjoint compact sets in the domain are mapped to compact sets which intersect in a set of measure zero.

One can actually obtain mappings $h : \mathbf{R}^r \rightarrow \mathbf{R}^s$ which satisfy (9.5) from limits of rescalings of g . This follows from Proposition 12.42 of [DS4]. These new mappings will also be Hölder continuous of the correct exponent and satisfy the regularity property discussed above. This measure-preserving property (9.5) is much closer to injectivity – see [DS4] for more information about it – but still it is not as strong as the bijectivity in Theorem 9.1.

It seems that no matter what we do in the continuous case we can never quite recapture fully the content of Theorem 9.1. Infinite processes are involved in both cases, but the condition of bijectivity in Theorem 9.1 does not appear to have a reasonable formulation in the infinite case without losing information, despite the fact that we can get a lot of structure for the mappings in the continuous case.

In neither case is the precise nature of the infinite processes required so clearly understood. One has examples and simple observations about what kind of behavior is not possible, but deeper results are missing.

9.2 A topological result about infinite complexity

By a *k-cell* we mean a topological space which is homeomorphic to the Cartesian product of k copies of the unit interval $[0, 1]$.

Theorem 9.6 ([Bry]) *Let D be a compact subset of some $\mathbf{R}^n$ which is homeomorphic to a cell of dimension k . Let $\mathbf{R}^n/D$ denote the topological space defined as a quotient of $\mathbf{R}^n$ by collapsing D to a point but leaving untouched the remaining elements of $\mathbf{R}^n$. Then $(\mathbf{R}^n/D) \times \mathbf{R}$ is homeomorphic to $\mathbf{R}^{n+1}$.*

Thus one might say that shrinking a cell to a point does not alter the ambient topology too painfully. If the cell is a standard one (a flat cube), then one can check directly that the quotient space $\mathbf{R}^n/D$ is homeomorphic to $\mathbf{R}^n$. There are however “wild” cells for which this is not the case. Theorem 9.6 is a bit remarkable for saying that after a single stabilization the problem always goes away.

This is a theorem about infinite complexity. The assumptions on D allow infinite processes, and so does the conclusion. It is a positive statement about controlling infinite complexity.

There are many other results of this general nature. See [Dm2] for more information.

9.3 Almost flat hypersurfaces

Let M be an hypersurface in $\mathbf{R}^{n+1}$. For simplicity we make the strong a priori assumptions that M is a smooth embedded submanifold which behaves well at infinity, e.g., is asymptotic to a hyperplane. We shall be interested in estimates which do not depend on these assumptions in a quantitative way.

Let $\epsilon > 0$ be given. We shall say that M is ϵ -flat if it satisfies the following two properties. The first asks that for each x and y on M we can find a rectifiable curve γ in M which connects them and which has length at most $(1 + \epsilon)|x - y|$. The second asks that if $x \in M$ and $r > 0$ are given, then

$$(9.7) \quad \left| \frac{|B(x, r) \cap M|}{\nu_n r^n} - 1 \right| \leq \epsilon.$$

Here $|B(x, r) \cap M|$ means the usual n -dimensional volume of the intersection of the ball $B(x, r)$ with M , while ν_n denotes the volume of the unit ball in $\mathbf{R}^n$.

If $\epsilon = 0$ then M must simply be an ordinary hyperplane in $\mathbf{R}^{n+1}$. If ϵ is small then these conditions imply that M is somehow pretty flat.

Question 9.8 *Fix a positive integer n . Is it true that for every $\eta > 0$ there is an $\epsilon > 0$ so that if M is an ϵ -flat hypersurface in $\mathbf{R}^{n+1}$, then there is a bijection $h : \mathbf{R}^n \rightarrow M$ which is $(1 + \eta)$ -bilipschitz, i.e.,*

$$(9.9) \quad (1 + \eta)^{-1}|x - y| \leq |h(x) - h(y)| \leq (1 + \eta)|x - y|$$

for all $x, y \in \mathbf{R}^n$?

It is easy to show that the converse to this is true. More precisely, if M does admit such a $(1 + \eta)$ -bilipschitz parameterization by $\mathbf{R}^n$, then M is ϵ -flat, where $\epsilon \rightarrow 0$ (uniformly) as $\eta \rightarrow 0$.

The a priori smoothness conditions here are convenient but not really the point. This question is really about the comparison between two different measurements of infinite complexity. Both ϵ -flatness and the $(1 + \eta)$ -bilipschitz allow infinite processes, infinite amounts of spiralling, for instance. They allow one to have bumps of small but definite size at infinitely many

scales and locations at once. But do they define practically the same condition, as in the question above?

This is not known. There are some partial results, about controlling the geometry of ϵ -flat hypersurfaces, and finding parameterizations which behave pretty well if not bilipschitzly, and about analysis of functions and operators on such surfaces. A general theme is that practically anything that one might know for hypersurfaces which admit $(1 + \eta)$ -bilipschitz parameterizations can be proved directly for ϵ -flat hypersurfaces, as long as one allows some movement in the parameters, and one asks questions in a reasonably simple language. (One should not simply reformulate the existence of a bilipschitz parameterization in other terms.)

In affect one is able to obtain a lot of results about infinite complexity in this case, even if the existence of good parameterizations is not known.

See [Se2, Se3, Se4, Se7, Se9, Se10] for more information about these and related questions.

9.4 Big pieces of bilipschitz mappings

Theorem 9.10 *Let Q denote the unit cube in $\mathbf{R}^n$, and suppose that $f : Q \rightarrow \mathbf{R}^n$ is Lipschitz with some constant L , i.e.,*

$$(9.11) \quad |f(x) - f(y)| \leq L|x - y|$$

for all $x, y \in Q$. Assume also that the image of f is of substantial size, i.e., $|f(Q)| \geq \delta$, where $|f(Q)|$ denotes the Lebesgue measure of $f(Q)$ and δ is some positive number. Then there is an $\epsilon > 0$, depending only on n, L , and δ , and a closed subset E of Q , such that f satisfies is actually bilipschitz on E , with

$$(9.12) \quad |f(x) - f(y)| \geq \epsilon|x - y|$$

for all $x, y \in E$.

Of course the existence of a substantial bilipschitz piece like this automatically ensures the existence of a lower bound on $|f(Q)|$. The fact that the converse holds, with uniform bounds, is quite remarkable.

This theorem was first proved in [Dv2] (see also [Dv3]). Another proof was given in [J1]. The latter proof was simpler and gave a stronger result, to the effect that the domain Q of f could be decomposed into a bounded

number of pieces, in such a way that the restriction of f to all but one of these pieces is bilipschitz with a uniform bound, while the image of the remaining piece under f has small Lebesgue measure. The more involved methods of [Dv2, Dv3] enjoy extra flexibility which enable them to be applied in other situations.

There are classical theorems in a somewhat similar spirit. One can decompose Q into a countable number of measurable subsets, on each of which f is either bilipschitz or has image of measure 0. See [Fe] for some results of this type, and for related facts concerning the “area” of Lipschitz mappings. These results permit one to conclude that the jacobian of f is bounded from below on a set of substantial measure. This is a weaker conclusion than the existence of a substantial bilipschitz piece, as in the theorem above. Indeed, one of the nice features of the theorem is that it lends itself well to discretization, and indeed the discrete versions contain essentially the same information as the continuous version.

Thus Theorem 9.10 and the more classical results which preceded it illustrate well the difference between quantitative bounds and the more qualitative considerations of countable unions and sets of measure zero. The latter live within a more symmetric language of infinite processes which is easier to use. In Theorem 9.10 we must break some of the symmetry of the language, but we then obtain a more concrete result which is combinatorial in essence. This fits well with the discussion of Chapter 3. For instance, a key point in the classical theory is that an arbitrary countable union of sets of measure zero also has measure zero. It is not at all clear what this means concretely, and one can certainly not say that arbitrary finite unions of sets of small measure also have small measure. To get quantitative bounds one must go much further inside the structure of the situation.

Additional progress in finding large bilipschitz pieces of Lipschitz mappings has been obtained in [JKV].

The story of large bilipschitz pieces should be compared with the one about ϵ -flat hypersurfaces and the question about bilipschitz parameterizations. It should obviously be easier to find bilipschitz pieces than global parameterizations, but that this is true in a practical way is less obvious at first glance. There is a lot of technology available now for finding large bilipschitz pieces of given spaces (additional references include [DS1, DS3]), and there is also now a reasonable collection of counterexamples concerning the existence of global bilipschitz and other controlled parameterizations [Se7, Se6].

One can ask similar questions in a broader geometric context, beyond spaces which have roughly Euclidean structure. See [DS4] for more information.

Bibliography

- [Ah] L. Ahlfors, *Complex Analysis*, McGraw-Hill.
- [Al] P. Alestalo, *Uniform domains of higher order*, Annales Academiæ Scientiarum Fennicæ Mathematica Ser. A I Math. Dissertationes **94** (1994), 1–48.
- [AV1] P. Alestalo and J. Väisälä, *Uniform domains of higher order II*, Annales Academiæ Scientiarum Fennicæ Mathematica Ser. A I Math. **21** (1996), 411–437.
- [AV2] P. Alestalo and J. Väisälä, *Uniform domains of higher order III*, Annales Academiæ Scientiarum Fennicæ Mathematica Ser. A I Math. **22** (1997), 445–464.
- [Ar] V. Arnold, *Mathematical Methods of Classical Mechanics*, Graduate Texts in Mathematics **60**, Springer-Verlag, 1989.
- [At1] O. Attie, *Quasi-isometry classification of some manifolds of bounded geometry*, Mathematische Zeitschrift **216** (1994), 501–527.
- [At2] O. Attie, *A surgery theory for manifolds of bounded geometry*, preprint.
- [AtH] O. Attie and S. Hurder, *Manifolds which cannot be leaves of foliations*, Topology **35**(1996), 335–353.
- [Ba] M. Barnesly, *Fractal image compression*, Notices of the American Mathematical Society **43** (1996), 657–662.
- [BCR1] G. Beylkin, R. Coifman, and V. Rokhlin, *Fast wavelet transforms and numerical algorithms I*, Communications on Pure and Applied Mathematics **44** (1991), 141–183.

- [BCR2] G. Beylkin, R. Coifman, and V. Rokhlin, *Wavelets in numerical analysis*, in *Wavelets and their applications* (M. Ruskai et al., eds.), Jones and Bartlett Publishers (1992), 181–210.
- [Bi] P. Billingsley, *Ergodic theory and information*, Wiley, 1965.
- [Br] G. Bredon, *Topology and Geometry*, Graduate Texts in Mathematics **139**, Springer-Verlag, 1993.
- [Bry] J. Bryant, *Euclidean space modulo a cell*, *Fundamenta Mathematicae* **63** (1968), 43–51.
- [BuC] C. Burgess and J. Cannon, *Embeddings of surfaces in E^3* , *Rocky Mountain Journal of Mathematics* **1** (1971), 259–344.
- [C1] J. Cannon, *Shrinking cell-like decompositions of manifolds: codimension 3*, *Annals of Mathematics* (2) **110** (1979), 83–112.
- [C2] J. Cannon, *The characterization of topological manifolds of dimension $n \geq 5$* , *Proceedings of the International Congress of Mathematicians* (Helsinki, 1978), 449–454.
- [Ca] A. Carbone, *Cycling in proofs and feasibility*, *Transactions of the American Mathematical Society* **352** (2000), 2049–2075.
- [CS1] A. Carbone and S. Semmes, *Making proofs without Modus Ponens: An introduction to the combinatorics and complexity of cut elimination*, *Bulletin of the American Mathematical Society* **34** (1997), 131–159.
- [CS2] A. Carbone and S. Semmes, *Propositional Proofs via Combinatorial Geometry and the Search for Symmetry*, in *Collegium Logicum, Annals of the Kurt-Gödel-Society, Volume 3*, 85–98, Institute of Computer Science AS CR, Prague, 1999.
- [CS3] A. Carbone and S. Semmes, *Looking from the inside and the outside*, *Synthese* **125** (2000), 385–416.
- [CS4] A. Carbone and S. Semmes, *A Graphic Apology for Symmetry and Implicitness*, Oxford University Press, 2000.

- [Ch] J. Cheeger, *Finiteness theorems for Riemannian manifolds*, American Journal of Mathematics **92** (1970), 61–74.
- [CK] J. Cheeger and J. Kister, *Counting topological manifolds*, Topology **9** (1970), 149–151.
- [Co] R. Coifman, *Adapted multiresolution analysis, computation, signal processing, and operator theory*, Proceedings of the International Congress of Mathematicians (Kyoto, 1990), Volume II , 879–887, 1991.
- [CMQW] R. Coifman, Y. Meyer, S. Quake, and V. Wickerhauser, *Signal processing and compression with wavelet packets*, in *Wavelets and their applications*, Proceedings of the NATO ASI Conference, 16–29 August, 1992 (J. Byrnes et al. eds.), Kluwer Academic Publishers, Nato ASI Series, Ser. C, Math. Phys. Sci. **442** (1994), 363–379.
- [CW] R. Coifman and G. Weiss, *Transference methods in analysis*, CBMS Lectures **31**, American Mathematical Society, 1977.
- [CW1] R. Coifman and V. Wickerhauser, *Wavelet analysis and signal processing*, in *Wavelets and Their Applications* (M. Ruskai et al., eds.), 153–178, Jones and Bartlett Publishers, 1992.
- [CW2] R. Coifman and V. Wickerhauser, *Wavelets and adapted waveform analysis. A toolkit for signal processing and numerical analysis*, in *Different perspectives on wavelets* (I. Daubechies, ed.), Proceedings of Symposia in Applied Mathematics **47**, 119–153, American Mathematical Society, 1993.
- [CR] S. Cook and R. Reckhow, *The relative efficiency of propositional proof systems*, Journal of Symbolic Logic **44** (1979), 36–50.
- [Db1] I. Daubechies, *Orthonormal bases of compactly supported wavelets*, Communications on Pure and Applied Mathematics **41** (1988), 901–996.
- [Db2] I. Daubechies, *Ten Lectures on Wavelets*, CBMS–NSF Regional conference series in Applied Mathematics **61**, Philadelphia, Pennsylvania, SIAM (Society for Industrial and Applied Mathematics), 1992.

- [Db3] I. Daubechies, *Wavelet transforms and orthonormal wavelet bases*, in *Different Perspectives on Wavelets* (I. Daubechies, ed.), Proceedings of Symposia in Applied Mathematics **47**, 1–33, American Mathematical Society, 1993.
- [Dm1] R. Daverman, *Embeddings of $(n-1)$ -spheres in Euclidean n -space*, Bulletin of the American Mathematical Society **84** (1978), 377–405.
- [Dm2] R. Daverman, *Decompositions of Manifolds*, Academic Press, 1986.
- [Dv1] G. David, *Opérateurs d'intégrale singulière sur les surfaces régulières*, Annales Scientifiques de l'École Normale Supérieure (4) **21** (1988), 225–258.
- [Dv2] G. David, *Morceaux de graphes lipschitziens et intégrales singulières sur un surface*, Revista Matemática Iberoamericana **4** (1988), 73–114.
- [Dv3] G. David, *Wavelets and Singular Integrals on Curves and Surfaces*, Lecture Notes in Mathematics **1465**, Springer-Verlag, 1991.
- [DS1] G. David and S. Semmes, *Singular Integrals and Rectifiable Sets in $\mathbf{R}^n$: Au-delà des graphes lipschitziens*, Astérisque **193**, Société Mathématique de France, 1991.
- [DS2] G. David and S. Semmes, *Harmonic Analysis and the geometry of subsets of $\mathbf{R}^n$* , Publications Mathématiques **35** (1991), 237–249.
- [DS3] G. David and S. Semmes, *Analysis of and on Uniformly Rectifiable Sets*, Mathematical Surveys and Monographs **38**, American Mathematical Society, 1993.
- [DS4] G. David and S. Semmes, *Fractured Fractals and Broken Dreams: Self-Similar Geometry through Metric and Measure*, Oxford University Press, 1997.
- [DoS] S. Donaldson and D. Sullivan, *Quasiconformal 4-manifolds*, Acta Mathematica **163** (1989), 181–252.
- [Do] J. Dorronsoro, *A characterization of potential spaces*, Proceedings of the American Mathematical Society **95** (1985), 21–31.

- [E] R. Edwards, *The topology of manifolds and cell-like maps*, Proceedings of the International Congress of Mathematicians (Helsinki, 1978), 111–127.
- [EFT] H.-D. Ebbinghaus, J. Flum, and W. Thomas, *Mathematical Logic*, Second Edition, Springer-Verlag, 1994.
- [Fa] K. Falconer, *The Geometry of Fractal Sets*, Cambridge University Press, 1984.
- [Fe] H. Federer, *Geometric Measure Theory*, Springer-Verlag, 1969.
- [F1] S. Ferry, *Homotoping ϵ -maps to homeomorphisms*, American Journal of Mathematics **101** (1979), 567–582.
- [F2] S. Ferry, *Topological finiteness theorems for manifolds in Gromov–Hausdorff space*, Duke Mathematical Journal **74** (1994), 95–106.
- [FM] B. Farb and L. Mosher, *A rigidity theorem for the solvable Baumslag–Solitar groups*, Inventiones Mathematicae **131** (1998), 419–451.
- [FQ] M. Freedman and F. Quinn, *Topology of 4-Manifolds*, Princeton University Press, 1990.
- [Ga] J. Garnett, *Bounded Analytic Functions*, Academic Press, 1981.
- [Gia] M. Giaquinta, *Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*, Annals of Mathematics Studies **105**, Princeton University Press, 1983.
- [Gir] J.-Y. Girard, *Proof Theory and Logical Complexity, Studies in Proof Theory 1*, Bibliopolis, Napoli, 1987.
- [GJ] M. Garey and D. Johnson, *Computers and Intractability*, W. H. Freeman and Company, 1979.
- [Gr1] M. Gromov, *Structures Métriques pour les Variétés Riemanniennes* (J. Lafontaine, P. Pansu, eds.), Cedic/Fernand Nathan, Paris, 1981. English translation in *Metric Structures for Riemannian and Non-Riemannian Spaces*, Birkhäuser, 1999.

- [Gr2] M. Gromov, *Groups of polynomial growth and expanding maps*, Publications Mathématiques (IHES) **53** (1981), 53–78.
- [Gr3] M. Gromov, *Hyperbolic groups*, in *Essays in Group Theory* (S. Gersten ed.), MSRI Publications **8**, p75–265, Springer-Verlag, 1987.
- [Gr4] M. Gromov, *Asymptotic invariants of infinite groups*, Proceedings of the 1991 conference Sussex conference on Geometric Group Theory (G. Niblo and M. Roller, eds.), London Mathematical Society Lecture Notes Series **182**, Cambridge University Press, 1993.
- [GP] M. Gromov and P. Pansu, *Rigidity of lattices: An introduction*, in *Geometric Topology: Recent Developments*, Lecture Notes in Mathematics **1504**, 39–137, Springer-Verlag, 1991.
- [GPW] K. Grove, P. Petersen V, and J.-Y. Wu, *Geometric finiteness theorems via controlled topology*, Inventiones Mathematicae **99** (1990), 205–213; erratum, **104** (1991), 221–222.
- [HSe] J. Heinonen and S. Semmes, *Thirty-three yes or no questions about mappings, measures, and metrics*, Conformal Geometry and Dynamics **1** (1997), 1–12 (an electronic journal of the American Mathematical Society).
- [HY] J. Heinonen and S. Yang, *Strongly uniform domains and periodic quasiconformal maps*, Annales Academiæ Scientiarum Fennicæ Mathematica Ser. A I Math. **20** (1995), 123–148.
- [HU] J. Hopcroft and J. Ullman, *Introduction to Automata Theory, Languages, and Computation*, Addison-Wesley, 1979.
- [HSt] E. Hewitt and K. Stromberg, *Real and Abstract Analysis*, Springer-Verlag, 1965.
- [HW] W. Hurewicz and H. Wallman, *Dimension Theory*, Princeton University Press, 1941, 1969.
- [J1] P. Jones, *Lipschitz and bi-Lipschitz functions*, Revista Matemática Iberoamericana **4** (1988), 115–122.
- [J2] P. Jones, *Rectifiable sets and the travelling salesman problem*, Inventiones Mathematicae **102** (1990), 1–15.

- [JKV] P. Jones, N. Katz, and A. Vargas, *Checkerboards, Lipschitz functions, and uniform rectifiability*, Revista Matemática Iberoamericana **13** (1997), 189–210.
- [Ka] Y. Katznelson, *An Introduction to Harmonic Analysis*, Dover Publications, 1976.
- [Ki] R. Kirby, *The Topology of 4-Manifolds*, Lecture Notes in Mathematics **1374**, Springer-Verlag, 1989.
- [KS] R. Kirby and L. Siebenmann, *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, Annals of Mathematics Studies **88**, Princeton University Press, 1977.
- [Man] Y. Manin, *A Course in Mathematical Logic*, Springer-Verlag, 1977.
- [Mat] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge University Press, 1995.
- [Mau] B. Maurey, *Théorèmes de factorisation pour les operateurs linéaires á valeurs dans les espaces L^p* , Astérisque **11**, Société Mathématique de France, 1974.
- [Me1] Y. Meyer, *Ondelettes et Opérateurs I: Ondelettes*, Hermann, 1990.
- [Me2] Y. Meyer, *Ondelettes et Opérateurs II: Opérateurs de Calderón-Zygmund*, Hermann, 1990.
- [Mi1] J. Milnor, *Two complexes which are homeomorphic but combinatorially distinct*, Annals of Mathematics (2) **74** (1961), 575–590.
- [Mi2] J. Milnor, *Topology from the Differentiable Viewpoint*, The University Press of Virginia, Charlottesville, 1965.
- [MiS] J. Milnor and J. Stasheff, *Characteristic Classes*, Annals of Mathematics Studies **76**, Princeton University Press, 1974.
- [Moi] E. Moise, *Geometric Topology in Dimensions 2 and 3*, Springer-Verlag, 1977.
- [Mor] F. Morgan, *Geometric Measure Theory: A Beginner's Guide*, Academic Press, 1988.

- [Mos] G. Mostow, *Strong Rigidity of Locally Symmetric Spaces*, Annals of Mathematics Studies **78**, Princeton University Press, 1973.
- [N] E. Nikishin, *Resonance theorems and superlinear operators*, Russian Mathematical Surveys **25** # 6 (1970), 125–187.
- [O] K. Okikiolu, *Characterization of subsets of rectifiable curves in $\mathbf{R}^n$* , Journal of the London Mathematical Society, Ser II **46** (1992), 336–348.
- [Pa] P. Pansu, *Métriques de Carnot–Carathéodory et quasiisométries des espaces symétriques de rang un*, Annals of Mathematics **129** (1989), 1–60.
- [Pap] C. Papdimitriou, *Computational Complexity*, Addison-Wesley, 1994.
- [Pe1] P. Petersen V, *A finiteness theorem for metric spaces*, Journal of Differential Geometry **31** (1990), 387–395.
- [Pe2] P. Petersen V, *Gromov–Hausdorff convergence of metric spaces*, Proceedings of Symposia in Pure Mathematics **54**, Part 3, 489–504, American Mathematical Society, 1993.
- [RS] R. Rochberg and S. Semmes, *Nearly weakly orthonormal sequences, singular value estimates, and Calderón–Zygmund operators*, Journal of Functional Analysis **86** (1989), 237–306.
- [Ru1] W. Rudin, *Principles of Mathematical Analysis*, McGraw-Hill, 1976.
- [Ru2] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, 1987.
- [Ru3] W. Rudin, *Functional Analysis*, McGraw-Hill, 1973.
- [Se1] S. Semmes, *Analysis vs. geometry on a class of rectifiable hypersurfaces in $\mathbf{R}^n$* , Indiana University Mathematics Journal **39** (1990), 1005–1035.
- [Se2] S. Semmes, *Chord–arc surfaces with small constant I* , Advances in Mathematics **85** (1991), 198–223.

- [Se3] S. Semmes, *Chord–arc surfaces with small constant II: Good parameterizations*, *Advances in Mathematics* **88** (1991), 170–199.
- [Se4] S. Semmes, *Hypersurfaces in $\mathbf{R}^n$ whose unit normal has small BMO norm*, *Proceedings of the American Mathematical Society* **112** (1991), 403–412.
- [Se5] S. Semmes, *Finding structure in sets with little smoothness*, *Proceedings of the International Congress of Mathematicians (Zürich, 1994)*, 875–885, Birkhäuser, 1995.
- [Se6] S. Semmes, *Good metric spaces without good parameterizations*, *Revista Matemática Iberoamericana* **12** (1996), 187–275.
- [Se7] S. Semmes, *On the nonexistence of bilipschitz parameterizations and geometric problems about A_∞ weights*, *Revista Matemática Iberoamericana* **12** (1996), 337–410.
- [Se8] S. Semmes, *Finding Curves on General Spaces through Quantitative Topology with Applications to Sobolev and Poincaré Inequalities*, *Selecta Mathematica (N.S.)* **2** (1996), 155–295.
- [Se9] S. Semmes, *Metric Spaces and Mappings Seen at Many Scales*, appendix in *Metric Structures for Riemannian and Non-Riemannian Spaces* by M. Gromov et al., Birkhäuser, 1999.
- [Se10] S. Semmes, *Mappings and Spaces*, in *Quasiconformal Mappings and Analysis: A Collection of Papers Honoring F.W. Gehring* (P. Duren, J. Heinonen, B. Osgood, and B. Palka, eds.), 347–368, Springer-Verlag, 1998.
- [Si] L. Siebenmann, *Topological Manifolds*, *Proceedings of the International Congress of Mathematicians (Nice, 1970)*, Volume 2, 133–163, Gauthier-Villars, Paris, 1971.
- [Smb] B. Simon, *Trace Ideals and their Applications*, *London Mathematical Society Lecture Note Series* **35**, Cambridge University Press, 1979.
- [Sml] L. Simon, *Lectures on Geometric Measure Theory*, *Proceedings of the Centre for Mathematical Analysis, Australian National University* **3**, 1984.

- [Sn] Y. Sinai, *Introduction to Ergodic Theory*, Mathematical Notes **18** (1976), Princeton University Press.
- [SS] L. Siebenmann and D. Sullivan, *On complexes that are Lipschitz manifolds*, in *Geometric Topology (Athens, Georgia, 1977)* (J. Cantrell, ed.), Academic Press, 1979, 503–525.
- [St1] E. Stein, *On limits of sequences of operators*, Annals of Mathematics (2) **74** (1961), 140–170.
- [St2] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press 1970.
- [SW] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [Sto] R. Stong, *Mapping $\mathbf{Z}^r$ into $\mathbf{Z}^s$ with maximal contraction*, Discrete and Computational Geometry **20** (1998), 131–138.
- [Su] D. Sullivan, *Hyperbolic geometry and homeomorphisms*, in *Geometric Topology (Athens, Georgia, 1977)* (J. Cantrell, ed.), Academic Press, 1979, 543–555.
- [T] R. Thom, *Quelques propriétés globales des variétés différentiables*, Commentarii Mathematici Helvetici **28** (1954), 17–86.
- [V] J. Väisälä, *Metric duality in Euclidean spaces*, Mathematica Scandinavica **80** (1997), 249–288.
- [W] F. Warner, *Foundations of Differentiable Manifolds and Lie Groups*, Scott, Foresman and Company, 1971.
- [Z] A. Zygmund, *Trigonometric Series*, Volumes I and II, Cambridge University Press, 1979.